

СИБИРСКИЕ ЭЛЕКТРОННЫЕ
МАТЕМАТИЧЕСКИЕ ИЗВЕСТИЯ

Siberian Electronic Mathematical Reports

<http://semr.math.nsc.ru>

Том 14, стр. 317–366 (2017)
DOI 10.17377/semi.2017.14.029

УДК 519.217.2, 519.681.2, 519.681.3
MSC 60J27, 60K15, 68Q85, 68Q87

EQUIVALENCES FOR FLUID STOCHASTIC PETRI NETS

I.V. TARASYUK, P. BUCHHOLZ

ABSTRACT. We propose fluid equivalences to compare and reduce behaviour of labeled fluid stochastic Petri nets (LFSPNs) while preserving their discrete and continuous properties. We define a linear-time relation of fluid trace equivalence and its branching-time counterpart, fluid bisimulation equivalence. Both fluid relations respect the essential features of the LFSPNs behaviour, such as *functional activity*, *stochastic timing* and *fluid flow*. We consider the LFSPNs whose continuous markings have no influence to the discrete ones, i.e. every discrete marking determines completely both the set of enabled transitions, their firing rates and the fluid flow rates of the incoming and outgoing arcs for each continuous place. We also require that the discrete part of the LFSPNs should be continuous time stochastic Petri nets. The underlying stochastic model for the discrete part of the LFSPNs is continuous time Markov chains (CTMCs). The performance analysis of the continuous part of LFSPNs is accomplished via the associated stochastic fluid models (SFMs). We show that fluid trace equivalence preserves average potential fluid change volume for the transition sequences of every certain length. We prove that fluid bisimulation equivalence preserves the following aggregated (by such a bisimulation) probability functions: stationary probability mass for the underlying CTMC, as well as stationary fluid buffer empty probability, fluid density and distribution for the associated SFM. Fluid bisimulation equivalence is then used to simplify the qualitative and quantitative analysis of LFSPNs that is accomplished by means of quotienting (by the equivalence) the discrete reachability graph and underlying CTMC. The application example of a document preparation system demonstrates the behavioural analysis via quotienting by fluid bisimulation equivalence.

TARASYUK I.V., BUCHHOLZ P., EQUIVALENCES FOR FLUID STOCHASTIC PETRI NETS.
© 2017 TARASYUK I.V., BUCHHOLZ P.

The work was partially supported by Deutsche Forschungsgemeinschaft (DFG) under grant BE 1267/14-1 and Russian Foundation for Basic Research (RFBR) under grant 14-01-91334.

Received December 7, 2016, published April 4, 2017.

Keywords: labeled fluid stochastic Petri net, continuous time stochastic Petri net, continuous time Markov chain, stochastic fluid model, transient and stationary behaviour, buffer empty probability, fluid density and distribution, performance analysis, Markovian trace and bisimulation equivalences, fluid trace and bisimulation equivalences, quotient, application.

1. INTRODUCTION

An important scientific problem that has been often addressed in the last decades is the design and analysis of parallel systems, which takes into account both qualitative (functional) and quantitative (timed, probabilistic, stochastic) features of their behaviour. The main goal of the research on this topic is the development of models and methods respecting performance requirements for concurrent and distributed systems with time constraints (such as deterministic, nondeterministic and stochastic time delays) to construct, validate and optimize the performability of realistic large-scale applications: computing systems, networks and software, controllers for industrial devices, manufacturing lines, vehicle, aircraft and transportation engines. A fruitful approach to achieving progress in this direction appeared to be a combined application of the theories of Petri nets, stochastic processes and fluid flow systems to the specification and analysis of such time-dependent systems with inherent behavioural randomness [35].

1.1. Fluid stochastic Petri nets. In the past, many extensions of stochastic Petri nets (SPNs) [48, 46, 47, 44, 45, 4, 5] have been developed to specify, model, simulate and analyze some particular classes of systems, such as computer systems, communication networks or manufacturing plants. These new formalisms have been constructed as a response to the needs for more expressive power in describing real-world systems, and to the requirements for compact models and efficient analysis techniques. One of the extensions are fluid stochastic Petri nets (FSPNs), capable of modeling hybrid systems that combine continuous state variables, corresponding to the fluid levels, with discrete state variables, specifying the token numbers. The continuous part of the FSPNs allows one to represent the fluid level in continuous places and fluid flow along continuous arcs. This part can naturally describe continuous variables in physical systems whose behaviour is commonly represented by differential equations. Continuous variables may also be used to describe a macroscopic view of discrete items that appear in large populations, e.g., packets in a computer network, molecules in a chemical reaction or people in a crowd. The discrete part of an FSPN is essentially its underlying SPN, obtained from the FSPN by removing all the fluid-related continuous elements. This part usually models the discrete control of the continuous process. The control may demonstrate some stochastic behavior that captures uncertainty about the detailed system behavior.

FSPNs have been proposed in [55, 22, 63] to model stochastic fluid flow systems [34, 30]. To analyze FSPNs, simulation, numerical and matrix-geometric methods are widely used [39, 23, 13, 31, 32, 28, 29, 40, 33]. The major problem of FSPNs is the high complexity of their solution, resulting in huge memory and time requirements while analyzing realistic models. A positive feature of the FSPN formalism is that it hides from a modeler the technical difficulties with solving differential equations for the underlying stochastic processes and that it unifies in one framework the evolution equations for the discrete and continuous parts of systems.

1.2. Equivalences on the related models. However, to the best of our knowledge, neither transition labeling nor behavioral equivalences have been proposed so far for FSPNs. In [56, 57, 58], label equivalence and projected label equivalence have been introduced for Fluid Process Algebra (FPA). FPA is a simple subalgebra of Grouped PEPA (GPEPA) [36], which is itself a conservative extension of Performance Evaluation Process Algebra (PEPA) [38], obtained by adding fluid semantics with an objective to simplify solving the systems of replicated ordinary differential equations. In [56, 58], it has been proved that projected label equivalence induces a fluid lumpable partition and that both label equivalence and projected label equivalence imply semi-isomorphism (stochastic isomorphism), in the context of a special subclass of well-posed models. Nevertheless, the mentioned label equivalences do not respect the action names; hence, they are not behavioral relations.

In [59, 60], the models specified with large ordinary differential equation (ODE) systems have been explored within Fluid Extended Process Algebra (FEPA). The relations of semi-isomorphism, ordinary and projected label equivalence have been proposed for the sequential process components, called fluid atoms, that can have a multiplicity (the number of copies in the model specification). In addition to exact fluid lumpability (EFL) from [56, 58] that aggregates isomorphic processes with the same multiplicities, ordinary fluid lumpability (OFL) has been proposed. OFL does not require that the multiplicities of the isomorphic processes coincide, but it preserves the sums of the aggregated variables instead. Moreover, the approximate versions (ϵ -variants) of semi-isomorphism, EFL and OFL have been investigated, which abstract from small fluctuations of the parameter values in the processes with close (similar) differential trajectories. However, the label equivalences do not respect the names of actions and therefore they are not behavioural equivalences.

In [61], two notions of lumpability for the class of heterogenous systems models specified by nonlinear ODEs have been investigated: exact lumpability (EL) [54] and uniform lumpability (UL), both applied for exact aggregation of the state variables. Unlike the EL transformations through linear mappings (in particular, those induced by a partition of the original state space), UL considers exact symmetries of the equations due to identification of the different variables from one partition block, which have coinciding differential trajectories (solutions) in case of the same initial conditions. This is an extension of the ODE systems reduction technique for *FPA* [56] to arbitrary vector fields. Both the lumpability relations do not take into account the action names and they are not behavioural equivalences.

In [41], differential bisimulation for FEPA has been constructed. This relation induces a partition on ODEs corresponding to the FEPA terms. Differential bisimulation is a behavioural equivalence that is an ODE analogue of the probabilistic and stochastic bisimulations. For each partition block, the sum of solutions of its ODEs coincides with the solution of a single aggregate ODE for this block. In the framework of FSPNs, the ODE systems are obtained only when there is exactly one continuous place. In the general case (more than one continuous place), the dynamics of FSPNs is described by the systems of equations with partial derivatives of probability distribution and density functions w.r.t. fluid levels in the continuous places. These levels are the random variables with a parameter accounting for the work time of an FSPN, starting from the initial moment. Just for the fluid levels, the ODEs over the time variable can be constructed in each discrete marking. However, the sojourn time in each discrete marking is a random variable, calculated

as the minimal transition delay, among all the transitions enabled in the marking. The FEPA processes are described by the ODE systems with derivatives of the population functions that define the multiplicities (numbers of replicas) of fluid atoms by only one variable denoting the time. Thus, the analogues of the FEPA fluid atoms are the (mainly, continuous) places of FSPNs. Hence, the FSPN model always has a naturally embedded notion of population, seen as a fluid in a continuous place. The systems behaviour is treated in FSPNs on a higher level of specification using the continuous time concept and the GSPN basic model, and also on a higher analysis level with constructing the underlying SMCs, CTMCs and stochastic fluid models (SFMs). The multiplicities of the FEPA fluid atoms are the functions of time, such that their values can be found for every particular time moment. In contrast, the fluid levels in continuous places of FSPNs are the continuous random variables that depend on time, so that their exact values at a given moment of time cannot be calculated. The reason is the property of the continuous probability distributions, stating that a continuous random variable may be equal to a concrete fixed value with zero probability only (excepting that in FSPNs, the fluid probability mass at the boundaries may be positive). In addition, the FEPA expressivity is restricted by considering only the processes, each being a parallel composition of the fluid atoms denoting a large number of copies of the simple sequential components, specified with only three operations: prefix, choice and recursive definition with constants. Moreover, the fluid atoms in FEPA are considered uniformly, i.e. there is no difference between “discrete” atoms with small multiplicities and “continuous” ones with large multiplicities. However, the tokens in FSPNs are jumped from one discrete place to another instantaneously when their input or output transitions fire, whereas the fluid flow proceeds through continuous places during all the time period when their input or output transitions are enabled. Thus, the notion of differential bisimulation cannot be straightforwardly transferred from FEPA to FSPNs.

In [17, 18, 20], back and forth bisimulation equivalences on chemical species have been introduced for chemical reaction networks (CRNs) with the ODE-based semantics. The forth bisimulation induces a partition where each equivalence class is a sum of concentrations of the species from this class, and this relation guarantees the ordinary fluid lumping on the ODEs of CRNs. The back bisimulation relates the species with the same ODE solutions at all time points, starting from the moment for which their equal initial conditions have been defined, and this relation characterizes the exact fluid lumping on the ODEs of CRNs. The bisimulations proposed in [17] differ from the equivalences from [56, 57, 58, 59, 60], since the former ones relate single variables whereas the latter ones relate the sets of variables, such that each of them represents the behaviour of some sequential process. The CRNs dynamics is described by ODEs with derivatives w.r.t. one variable (time), and the CRNs behaviour is deterministic, described by differential trajectories. Unlike CRNs, FSPNs have a stochastic behaviour which is influenced by the interplay of time and probabilistic factors. The FSPNs dynamics is analyzed with SFMs, solved using the differential equations with partial derivatives w.r.t. several variables.

In [19], back and forth differential equivalences have been explored for a basic formalism, called Intermediate Drift Oriented Language (IDOL). IDOL has a syntax to specify drift for a class of non-linear ODEs, for which the decidability results are known. The mentioned equivalence relations can be transferred from IDOL to the higher-level models, such as Petri nets, process algebras and rule systems,

interpreted as ODEs. The differential equivalences embrace such notions as minimization of CTMCs based on the lumpability relation [26], bisimulations of CRNs [17] and behavioural relations for process algebras with the ODE semantics [41]. At the same time, the ODE class defined by the IDOL language cannot specify semantics of the systems with stochastic continuous time delays in the discrete states, as well as many other behavioural aspects of FSPNs, including the ones mentioned above.

In [1], on the product form queueing networks, the ideas of equivalent flow server and flow equivalence have been applied to the models reduction, by aggregating server stations and their states by that equivalence. Nevertheless, flow equivalence does not respect the names of actions, hence, it is not a behavioural relation.

1.3. Our contributions. In this paper, we propose the behavioural relations of fluid trace and bisimulation equivalences that are useful for the comparison and reduction of the behaviour of LFSPNs, since these relations preserve the functionality and performability of their discrete and continuous parts.

For every FSPN, the discrete part of its marking is determined by the natural number of tokens contained in the discrete places. The continuous places of an FSPN are associated with the non-negative real-valued fluid levels that determine the continuous part of the FSPN marking. Thus, FSPNs have a hybrid (discrete-continuous) state space. The discrete part of a hybrid marking of FSPNs is called discrete marking while the continuous part is called continuous one. The discrete part of each hybrid marking has an influence on the continuous part. For more general FSPNs, the reverse dependence is possible as well. As a basic model for constructing LFSPNs, we consider only the FSPNs in which the continuous parts of markings have no influence on the discrete ones, i.e. every discrete part determines completely both the set of enabled transitions and the rates of incoming and outgoing arcs for each continuous place [28, 33]. We also require that the discrete part of LFSPNs should be labeled continuous time stochastic Petri nets (CTSPNs) [46, 44, 45, 4].

First, we define a linear-time relation of *fluid trace equivalence* on LFSPNs. Linear-time equivalences, unlike branching-time ones, do not respect the points of choice among several alternative continuations of the system's behavior. We require the relation on discrete markings of two LFSPNs to be a standard (strong) Markovian trace equivalence. Hence, for every sequence of discrete markings and transitions in the discrete reachability graph of an LFSPN, starting from the initial discrete marking (such sequence is called path), we require a simulation of the path in the discrete reachability graph of the equivalent LFSPN, such that the action labels of the corresponding fired transitions in the both sequences coincide. Next, the average sojourn times in the respective discrete markings should be the same. Finally, for the two equivalent LFSPNs, the cumulative execution probabilities of all the paths corresponding to a particular sequence of actions, together with a concrete sequence of the average sojourn times, should be equal. Thus, when comparing the execution probabilities, we parameterize the paths with the same extracted action sequence by all possible sequences of the extracted average sojourn times. Our definition of the trace equivalence on the discrete markings of LFSPNs is similar to that of ordinary (that with the absolute time counter or with the countdown timer) Markovian trace equivalence [62] on transition-labeled CTMCs. Ordinary Markovian trace equivalence and its variants from [62] have been later investigated and enhanced on sequential and concurrent Markovian process calculi SMPC and CMPC in [8, 6, 7, 9] and on Uniform Labeled Transition Systems (ULTraS) in

[10, 11]. As for the continuous markings of the two LFSPNs, we further parameterize the paths with the same extracted action sequence and the same sequence of the extracted average sojourn times by counting the execution probabilities only of those paths additionally having the same sequence of extracted fluid flow rates of the respective continuous places in the corresponding discrete markings. We show that fluid trace equivalence preserves average potential fluid change volume in the respective continuous places for the transition sequences of each particular length.

Second, we propose a branching-time relation of *fluid bisimulation equivalence* on LFSPNs that is strictly stronger than fluid trace equivalence. We require the fluid bisimulation on the discrete markings of two LFSPNs to be a standard (strong) Markovian bisimulation. Hence, for each transition firing in an LFSPN, we require a simulation of the firing in the equivalent LFSPN, such that the action labels of the both fired transitions and their overall rates coincide. Thus, our definition of the bisimulation equivalence on the discrete markings of LFSPNs is similar to that of the performance bisimulation equivalences [15, 16] on labeled CTSPNs and labeled generalized SPNs (GSPNs) [44, 21, 45, 14, 4, 5], as well as the strong equivalence from [38] on stochastic process algebra PEPA. All these relations belong to the family of Markovian bisimulation equivalences, investigated on sequential and concurrent Markovian process calculi SMPC and CMPC in [8, 6, 7, 9], as well as on Uniform Labeled Transition Systems (ULTraS) in [10, 11]. As for the continuous markings, we should fix a bijective correspondence between the sets of continuous places of the two LFSPNs. We require that, for every pair of the Markovian bisimilar discrete markings, the fluid flow rates of the continuous places in the first LFSPN should coincide with those of the corresponding continuous places in the second LFSPN. We prove that fluid bisimulation equivalence preserves, for the equivalence classes, the stationary probability distribution of the underlying continuous time Markov chain (CTMC), as well as the stationary fluid buffer empty probability, probability distribution and density for the associated stochastic fluid model (SFM). Hence, the equivalence guarantees identity of many performance measures, calculated for the stationary behaviour of the LFSPNs. The fluid bisimulation equivalence is also used to simplify the analysis of LFSPNs, due to diminishing the number of discrete markings that are lumped into the equivalence classes, interpreted as the states of the quotient discrete reachability graph and quotient underlying CTMC. The quotients of the probability functions describe the quotient associated SFM.

The application example of the three LFSPNs modeling the document preparation system demonstrates how the LFSPNs structure and behaviour can be reduced by fluid bisimulation equivalence while preserving their behavioural properties.

The first results on this subject can be found in [53], where we have proposed a class of LFSPNs and defined a novel fluid bisimulation equivalence for them that preserves aggregate fluid density and distribution, as well as discrete and continuous performance measures. This paper extends that publication with the new results for LFSPNs: fluid trace equivalence, interrelations of the fluid equivalences, quotienting by fluid bisimulation, the probability functions quotients and application example.

1.4. Outline of the paper. In Section 2, we present the definition and behaviour of LFSPNs. Section 3 explores the discrete part of LFSPNs, i.e. the derived labeled CTSPNs and their underlying CTMCs. Section 4 investigates the continuous part of LFSPNs, which is the associated SFMs. In Section 5, we construct a linear-time

relation of fluid trace equivalence for LFSPNs. In Section 6, we propose a branching-time relation of fluid bisimulation equivalence for LFSPNs and compare it with the fluid trace one. In Section 7, we explain how to reduce discrete reachability graphs and underlying CTMCs of LFSPNs modulo fluid bisimulation equivalence, by applying quotienting. Section 8 contains the preservation results for the quantitative behaviour of LFSPNs modulo fluid bisimulation equivalence. Section 9 describes a case study of three LFSPNs modeling the document preparation system. Section 10 summarizes the results obtained and outlines research perspectives in this area.

2. BASIC CONCEPTS OF LFSPNS

Let us introduce a class of labeled fluid stochastic Petri nets (LFSPNs), whose transitions are labeled with action names, used to specify different system activities. Without labels, LFSPNs are essentially a subclass of FSPNs [39, 28, 33], so that their discrete part describes CTSPNs [46, 44, 45, 4]. This means that LFSPNs have no inhibitor arcs, priorities and immediate transitions, which are used in the standard FSPNs, which are the continuous extension of GSPNs. However, in many practical applications, the performance analysis of GSPNs is simplified by transforming them into CTSPNs or reducing their underlying semi-Markov chains into CTMCs (which are the underlying stochastic process of CTSPNs) by eliminating vanishing states [21, 45, 4, 5]. Transition labeling in LFSPNs is similar to the labeling, proposed for CTSPNs in [15]. Moreover, we suppose that the firing rates of transitions and flow rates of the continuous arcs do not depend on the continuous markings (fluid levels).

Let $\mathbb{N} = \{0, 1, 2, \dots\}$ be the set of *all natural numbers* and $\mathbb{N}_{\geq 1} = \{1, 2, \dots\}$ be the set of *all positive natural numbers*. Further, let $\mathbb{R} = (-\infty; \infty)$ be the set of *all real numbers*, $\mathbb{R}_{\geq 0} = [0; \infty)$ be the set of *all non-negative real numbers* and $\mathbb{R}_{> 0} = (0; \infty)$ be the set of *all positive real numbers*. The set of *all row vectors of $n \in \mathbb{N}_{\geq 1}$ elements from a set X* is defined as $X^n = \{(x_1, \dots, x_n) \mid x_i \in X \ (1 \leq i \leq n)\}$. The set of *all mappings from a set X to a set Y* is defined as $Y^X = \{f \mid f : X \rightarrow Y\}$. Let $Act = \{a, b, \dots\}$ be the set of *actions*.

First, we present a formal definition of LFSPNs.

Definition 1. A labeled fluid stochastic Petri net (LFSPN) is a tuple $N = (P_N, T_N, W_N, C_N, R_N, \Omega_N, L_N, \mathcal{M}_N)$, where

- $P_N = Pd_N \uplus Pc_N$ is a finite set of discrete and continuous places (\uplus denotes disjoint union);
- T_N is a finite set of transitions, such that $P_N \cup T_N \neq \emptyset$ and $P_N \cap T_N = \emptyset$;
- $W_N : (Pd_N \times T_N) \cup (T_N \times Pd_N) \rightarrow \mathbb{N}$ is a function providing the weights of discrete arcs between discrete places and transitions;
- $C_N \subseteq (Pc_N \times T_N) \cup (T_N \times Pc_N)$ is the set of continuous arcs between continuous places and transitions;
- $R_N : C_N \times \mathbb{N}^{|Pd_N|} \rightarrow \mathbb{R}_{\geq 0}$ is a function providing the flow rates of continuous arcs in a given discrete marking (the markings will be defined later);
- $\Omega_N : T_N \times \mathbb{N}^{|Pd_N|} \rightarrow \mathbb{R}_{> 0}$ is the transition rate function associating transitions with rates in a given discrete marking;
- $L_N : T_N \rightarrow Act$ is the transition labeling function assigning actions to transitions;
- $\mathcal{M}_N = (M_N, \mathbf{0})$, where $M_N \in \mathbb{N}^{|Pd_N|}$ and $\mathbf{0}$ is a row vector of $|Pc_N|$ values 0, is the initial (discrete-continuous) marking.

Let us consider in more detail the tuple elements from the definition above.

Every discrete place $p_i \in Pd_N$ may contain discrete tokens, whose number is represented by a natural number $M_i \in \mathbb{N}$ ($1 \leq i \leq |Pd_N|$). Each continuous place $q_j \in Pc_N$ may contain continuous fluid, with the level represented by a non-negative real number $X_j \in \mathbb{R}_{\geq 0}$ ($1 \leq j \leq |Pc_N|$). Then the complete hybrid (discrete-continuous) marking of N is a pair (M, X) , where $M = (M_1, \dots, M_{|Pd_N|})$ is a discrete marking and $X = (X_1, \dots, X_{|Pc_N|})$ is a continuous marking. When needed, these vectors can also be seen as the mappings $M : Pd_N \rightarrow \mathbb{N}$ with $M(p_i) = M_i$ ($1 \leq i \leq |Pd_N|$) and $X : Pc_N \rightarrow \mathbb{R}_{\geq 0}$ with $X(q_j) = X_j$ ($1 \leq j \leq |Pc_N|$). The set of *all markings (reachability set)* of N is denoted by $RS(N)$. Then $DRS(N) = \{M \mid (M, X) \in RS(N)\}$ is the set of *all discrete markings (discrete reachability set)* of N . $DRS(N)$ will be formally defined later. Further, $CRS(N) = \{X \mid (M, X) \in RS(N)\} \subseteq \mathbb{R}_{\geq 0}^{|Pc_N|}$ is the set of *all continuous markings (continuous reachability set)* of N . Every marking $(M, X) \in RS(N)$ evolves in time, hence, we can interpret it as a stochastic process $\{(M(\delta), X(\delta)) \mid \delta \geq 0\}$. Then the initial marking of N is that at the zero time moment, i.e. $\mathcal{M}_N = (M_N, \mathbf{0}) = (M(0), X(0))$, where $X(0) = \mathbf{0}$ means that all the continuous places are initially empty.

Every transition $t \in T_N$ has an associated positive real-valued instantaneous rate $\Omega_N(t, M) \in \mathbb{R}_{> 0}$, which is a parameter of the exponential distribution governing the transition delay (being a random variable) in the current discrete marking M . Transitions are labeled with actions, each representing a sort of activity they model.

Every discrete arc $da = (p, t)$ or $da = (t, p)$, where $p \in Pd_N$ and $t \in T_N$, connects discrete places and transitions. It has a non-negative integer-valued weight $W_N(da) \in \mathbb{N}$ assigned, representing its multiplicity. The zero weight indicates that the corresponding discrete arc does not exist, since its multiplicity is zero in this case. In the discrete marking $M \in DRS(N)$, every continuous arc $ca = (q, t)$ or $ca = (t, q)$, where $q \in Pc_N$ and $t \in T_N$, connects continuous places and transitions. It has a non-negative real-valued flow rate $R_N(ca, M) \in \mathbb{R}_{\geq 0}$ of fluid through ca , when the current discrete marking is M . The zero flow rate indicates that the fluid flow along the corresponding continuous arc is stopped in some discrete marking.

The graphical representation of LFSPNs resembles that for standard labeled Petri nets, but supplemented with the rates or weights, written near the corresponding transitions or arcs. Discrete places are drawn with ordinary circles while double concentric circles correspond to the continuous ones. Square boxes with the action names inside depict transitions and their labels. Discrete arcs are drawn as thin lines with arrows at the end while continuous arcs should represent pipes, so the latter are depicted by thick arrowed lines. If the rates are not given in the picture then they are assumed to be of no importance in the corresponding examples. The names of places and transitions are depicted near them when needed.

We now consider the behaviour of LFSPNs.

Let N be an LFSPN and M be a discrete marking of N . A transition $t \in T_N$ is *enabled* in M if $\forall p \in Pd_N W_N(p, t) \leq M(p)$. Let $Ena(M)$ be the set of *all transitions enabled in M* . Firings of transitions are atomic operations, and only single transitions are fired at once. Note that the enabling condition depends only on the discrete part of N and this condition is the same as for CTSPNs. Firing of a transition $t \in Ena(M)$ changes M to another discrete marking \widetilde{M} , such as $\forall p \in Pd_N \widetilde{M}(p) = M(p) - W_N(p, t) + W_N(t, p)$, denoted by $M \xrightarrow{t} \widetilde{M}$, where $\lambda = \Omega_N(t, M)$. We write $M \xrightarrow{t} \widetilde{M}$ if $\exists \lambda M \xrightarrow{\lambda} \widetilde{M}$ and $M \rightarrow \widetilde{M}$ if $\exists t M \xrightarrow{t} \widetilde{M}$.

Let us formally define the discrete reachability set of N .

Definition 2. Let N be an LFSPN. The discrete reachability set of N , denoted by $DRS(N)$, is the minimal set of discrete markings such that

- $M_N \in DRS(N)$;
- if $M \in DRS(N)$ and $M \rightarrow \widetilde{M}$ then $\widetilde{M} \in DRS(N)$.

Let us now define the discrete reachability graph of N .

Definition 3. Let N be an LFSPN. The discrete reachability graph of N is a labeled transition system $DRG(N) = (S_N, \mathcal{L}_N, \mathcal{T}_N, s_N)$, where

- the set of states is $S_N = DRS(N)$;
- the set of labels is $\mathcal{L}_N = T_N \times \mathbb{R}_{>0}$;
- the set of transitions is $\mathcal{T}_N = \{(M, (t, \lambda), \widetilde{M}) \mid M, \widetilde{M} \in DRS(N), M \xrightarrow{t, \lambda} \widetilde{M}\}$;
- the initial state is $s_N = M_N$.

3. DISCRETE PART OF LFSPNs

We have restricted the class of FSPNs underlying LFSPNs to those whose discrete part is CTSPNs, since the performance analysis of standard FSPNs with GSPNs as the discrete part is finally based on the CTMCs which are extracted from the underlying semi-Markov chains (SMCs) of the GSPNs by removing vanishing states. Consider the behaviour of the discrete part of LFSPNs, which is labeled CTSPNs.

For an LFSPN N , a continuous random variable $\xi(M)$ is associated with every discrete marking $M \in DRS(N)$. The variable captures a residence (sojourn) time in M . We adopt the *race* semantics, in which the fastest stochastic transition (i.e. that with the minimal exponentially distributed firing delay) fires first. Hence, the *probability distribution function (PDF)* of the sojourn time in M is that of the minimal firing delay of transitions from $Ena(M)$. Since exponential distributions are closed under minimum, the sojourn time in M is (again) exponentially distributed with a parameter, called the *exit rate from the discrete marking M* , defined as

$$RE(M) = \sum_{t \in Ena(M)} \Omega_N(t, M).$$

Note that we may have $RE(M) = 0$, meaning that there is no exit from M , if it is a *terminal discrete marking*, i.e. there are no transitions from it to different ones.

Hence, the PDF of the sojourn time in M (the probability of the residence time in M being less than δ) is $F_{\xi(M)}(\delta) = \mathbb{P}(\xi(M) < \delta) = 1 - e^{-RE(M)\delta}$ ($\delta \geq 0$). Then the *probability density function* of the residence time in M (the limit probability of staying in M at the time δ) is $f_{\xi(M)}(\delta) = \lim_{\Delta \rightarrow 0} \frac{F_{\xi(M)}(\delta + \Delta) - F_{\xi(M)}(\delta)}{\Delta} = \frac{dF_{\xi(M)}(\delta)}{d\delta} = RE(M)e^{-RE(M)\delta}$ ($\delta \geq 0$). The mean value (average, expectation) formula for the exponential distribution allows us to calculate the average sojourn time in M as $\mathbb{M}(\xi(M)) = \int_0^\infty \delta f_{\xi(M)}(\delta) d\delta = \frac{1}{RE(M)}$. The variance (dispersion) formula for the exponential distribution allows us to calculate the sojourn time variance in M as $\mathbb{D}(\xi(M)) = \int_0^\infty (\delta - \mathbb{M}(\xi(M)))^2 f_{\xi(M)}(\delta) d\delta = \frac{1}{(RE(M))^2}$.

The *average sojourn time in the discrete marking M* is

$$SJ(M) = \frac{1}{\sum_{t \in Ena(M)} \Omega_N(t, M)} = \frac{1}{RE(M)}.$$

The *average sojourn time vector* of N , denoted by SJ , has the elements $SJ(M)$, $M \in DRS(N)$.

Note that we may have $SJ(M) = \infty$, meaning that we stay in M forever, if it is a terminal discrete marking.

The *sojourn time variance in the discrete marking* M is

$$VAR(M) = \frac{1}{\left(\sum_{t \in E_{na}(M)} \Omega_N(t, M)\right)^2} = \frac{1}{RE(M)^2}.$$

The *sojourn time variance vector* of N , denoted by VAR , has the elements $VAR(M)$, $M \in DRS(N)$.

Note that we may have $VAR(M) = \infty$, meaning that the variance of the infinite sojourn time in M is infinite too, if it is a terminal discrete marking.

To evaluate performance with the use of the discrete part of N , we should investigate the stochastic process associated with it. The process is the underlying continuous time Markov chain, denoted by $CTMC(N)$.

Let $M, \widetilde{M} \in DRS(N)$. The *rate of moving from M to \widetilde{M} by firing any transition* is

$$RM(M, \widetilde{M}) = \sum_{\{t | M \xrightarrow{t} \widetilde{M}\}} \Omega_N(t, M).$$

Definition 4. Let N be an LFSPN. The underlying continuous time Markov chain (CTMC) of N , denoted by $CTMC(N)$, has the state space $DRS(N)$, the initial state M_N and the transitions $M \rightarrow_\lambda \widetilde{M}$, if $M \rightarrow \widetilde{M}$, where $\lambda = RM(M, \widetilde{M})$.

Isomorphism is a coincidence of systems up to renaming their components or states. Let \simeq denote isomorphism between CTMCs that binds their initial states.

Let N be an LFSPN. The elements \mathcal{Q}_{ij} ($1 \leq i, j \leq n = |DRS(N)|$) of the *transition rate matrix (TRM)*, called *infinitesimal generator*, \mathbf{Q} for $CTMC(N)$ are

$$\mathcal{Q}_{ij} = \begin{cases} RM(M_i, M_j), & i \neq j; \\ -\sum_{\{k | 1 \leq k \leq n, k \neq i\}} RM(M_i, M_k), & i = j. \end{cases}$$

The transient *probability mass function (PMF)* $\varphi(\delta) = (\varphi_1(\delta), \dots, \varphi_n(\delta))$ for $CTMC(N)$ is calculated via matrix exponent as

$$\varphi(\delta) = \varphi(0)e^{\mathbf{Q}\delta},$$

where $\varphi(0) = (\varphi_1(0), \dots, \varphi_n(0))$ is the initial PMF, defined as

$$\varphi_i(0) = \begin{cases} 1, & M_i = M_N; \\ 0, & \text{otherwise.} \end{cases}$$

The steady-state PMF $\varphi = (\varphi_1, \dots, \varphi_n)$ for $CTMC(N)$ is a solution of the linear equation system

$$\begin{cases} \varphi \mathbf{Q} = \mathbf{0} \\ \varphi \mathbf{1}^T = 1 \end{cases},$$

where $\mathbf{0}$ is a row vector of n values 0 and $\mathbf{1}$ is that of n values 1.

Note that the vector φ exists and is unique, if $CTMC(N)$ is ergodic. Then $CTMC(N)$ has a single steady state, and we have $\varphi = \lim_{\delta \rightarrow \infty} \varphi(\delta)$.

Let N be an LFSPN. In [53], we have presented a number of the steady-state *discrete performance indices (measures)*, which can be calculated based on the steady-state PMF φ for $CTMC(N)$ [46, 44, 21, 14, 45, 4, 5].

4. CONTINUOUS PART OF LFSPNS

We now consider the impact the discrete part of LFSPNs has on their continuous part, which is stochastic fluid models (SFMs). We investigate LFSPNs with a single continuous place, since the definitions and our subsequent results on the fluid equivalences can be transferred straightforwardly to the case of several continuous places, where multidimensional SFMs have to be explored.

Let N be an LFSPN such that $P_{C_N} = \{q\}$ and $M(\delta) \in DRS(N)$ be its discrete marking at the time $\delta \geq 0$. Every continuous arc $ca = (q, t)$ or $ca = (t, q)$, where $t \in T_N$, changes the fluid level in the continuous place q at the time δ with the flow rate $R_N(ca, M(\delta))$. So, in the discrete marking $M(\delta)$ fluid can leave q along the continuous arc (q, t) with the rate $R_N((q, t), M(\delta))$ and can enter q along the continuous arc (t, q) with the rate $R_N((t, q), M(\delta))$ for every transition $t \in E_{na}(M(\delta))$.

The *potential rate of the fluid level change (fluid flow rate) for the continuous place q* in the discrete marking $M(\delta)$ is

$$RP(M(\delta)) = \sum_{\{t \in E_{na}(M(\delta)) | (t, q) \in C_N\}} R_N((t, q), M(\delta)) - \sum_{\{t \in E_{na}(M(\delta)) | (q, t) \in C_N\}} R_N((q, t), M(\delta)).$$

Let $X(\delta)$ be the fluid level in q at the time δ . It is clear that the fluid level in a continuous place can never be negative. Therefore, $X(\delta)$ satisfies the following ordinary differential equation describing the *actual fluid flow rate for the continuous place q* in the marking $(M(\delta), X(\delta))$:

$$RA(M(\delta), X(\delta)) = \frac{dX(\delta)}{d\delta} = \begin{cases} \max\{RP(M(\delta)), 0\}, & X(\delta) = 0; \\ RP(M(\delta)), & (X(\delta) > 0) \wedge \\ & (RP(M(\delta^-))RP(M(\delta^+)) \geq 0); \\ 0, & (X(\delta) > 0) \wedge \\ & (RP(M(\delta^-))RP(M(\delta^+)) < 0). \end{cases}$$

In the first case considered in the definition above, we have $X(\delta) = 0$. In this case, if $RP(M(\delta)) \geq 0$ then the fluid level is growing and the derivative is equal to the potential rate. Otherwise, if $RP(M(\delta)) < 0$ then we should prevent the fluid level from crossing the lower boundary (zero) by stopping the fluid flow. For an explanation of the more complex second and third cases please refer to [27, 39, 28, 33]. Note that $\frac{dX(\delta)}{d\delta}$ is a piecewise constant function of $X(\delta)$ during the time periods when $M(\delta)$ remains unchanged. Hence, for each different ‘‘constant’’ segment we have $\frac{dX(\delta)}{d\delta} = RP(M(\delta))$ or $\frac{dX(\delta)}{d\delta} = 0$ and, therefore, we can suppose that within each such segment $RP(M(\delta))$ or 0 are the *actual fluid flow rates* for the continuous place q in the marking $(M(\delta), X(\delta))$. While constructing differential equations that describe the behaviour of SFMs associated with LFSPNs, we are interested only in the segments where $\frac{dX(\delta)}{d\delta} = RP(M(\delta))$. The SFMs behaviour within the remaining segments, where $\frac{dX(\delta)}{d\delta} = 0$, is completely described by the buffer empty probability function that collects the probability mass at the lower boundary.

The elements \mathcal{R}_{ij} ($1 \leq i, j \leq n = |DRS(N)|$) of the *fluid rate matrix (FRM)* \mathbf{R} for the continuous place q are defined as

$$\mathcal{R}_{ij} = \begin{cases} RP(M_i), & i = j; \\ 0, & i \neq j. \end{cases}$$

According to [28, 33], the underlying SFM of LFSPNs is the first order, infinite buffer, homogeneous Markov fluid model. The discrete part of the SFM derived from an LFSPN N is the CTMC $CTMC(N)$ with the TRM \mathbf{Q} . The evolution of the continuous part of the SFM (the fluid flow drift) is described by the FRM \mathbf{R} .

Let us consider the *transient behaviour* of the SFM associated with an LFSPN N . We introduce the following transient probability functions.

- $\varphi_i(\delta) = \mathbb{P}(M(\delta) = M_i)$ is the *discrete marking probability*;
- $\ell_i(\delta) = \mathbb{P}(X(\delta) = 0, M(\delta) = M_i)$ is the *buffer empty probability (probability mass at the lower boundary)*;
- $F_i(\delta, x) = \mathbb{P}(X(\delta) < x, M(\delta) = M_i)$ is the *fluid probability distribution function*;
- $f_i(\delta, x) = \frac{\partial F_i(\delta, x)}{\partial x} = \lim_{h \rightarrow 0} \frac{F_i(\delta, x+h) - F_i(\delta, x)}{h} = \lim_{h \rightarrow 0} \frac{\mathbb{P}(x < X(\delta) < x+h, M(\delta) = M_i)}{h}$ is the *fluid probability density function*.

The initial conditions are:

$$\ell_i(0) = \begin{cases} 1, & M_i = M_N; \\ 0, & \text{otherwise}; \end{cases}$$

$$F_i(0, x) = \begin{cases} 1, & (M_i = M_N) \wedge (x \geq 0); \\ 0, & \text{otherwise}; \end{cases}$$

$$f_i(0, x) = 0 \quad \forall (M_i, x) \in RS(N).$$

Let $\varphi(\delta), \ell(\delta), F(\delta, x), f(\delta, x)$ be the row vectors with the elements $\varphi_i(\delta), \ell_i(\delta), F_i(\delta, x), f_i(\delta, x)$, respectively ($1 \leq i \leq n$).

By the total probability law, we have

$$\ell(\delta) + \int_{0+}^{\infty} f(\delta, x) dx = \varphi(\delta).$$

The partial differential equations describing the transient behaviour are

$$\frac{\partial F(\delta, x)}{\partial \delta} + \frac{\partial F(\delta, x)}{\partial x} \mathbf{R} = F(\delta, x) \mathbf{Q}, \quad x > 0;$$

$$\frac{\partial f(\delta, x)}{\partial \delta} + \frac{\partial f(\delta, x)}{\partial x} \mathbf{R} = f(\delta, x) \mathbf{Q}, \quad x > 0.$$

Note that we have $\frac{\partial F(\delta, x)}{\partial x} = f(\delta, x)$, $F(\delta, 0) = \ell(\delta)$, $F(\delta, \infty) = \varphi(\delta)$.

The partial differential equation for the buffer empty probabilities (lower boundary conditions) are

$$\frac{d\ell(\delta)}{d\delta} + f(\delta, 0) \mathbf{R} = \ell(\delta) \mathbf{Q}.$$

The lower boundary constraint is: if $\mathcal{R}_{ii} = RP(M_i) > 0$ then $\ell_i(\delta) = F_i(\delta, 0) = 0$ ($1 \leq i \leq n$).

The normalizing condition is

$$\ell(\delta)\mathbf{1}^T + \int_{0+}^{\infty} f(\delta, x)dx\mathbf{1}^T = 1,$$

where $\mathbf{1}$ is a row vector of n values 1.

Let us now consider the *stationary behaviour* of the SFM associated with an LFSPN N . We do not discuss here in detail the conditions under which the steady state for the associated SFM *exists* and is *unique*, since this topic has been extensively explored in [39, 28, 33]. Particularly, according to [39, 33], the steady-state PDF *exists* (i.e. the transient functions approach their stationary values, as the time parameter δ tends to infinity in the transient equations), when the associated SFM is a Markov fluid model, whose fluid flow drift (described by the matrix \mathbf{R}) and transition rates (described by the matrix \mathbf{Q}) are fluid level independent, and the following *stability condition* holds:

$$\text{FluidFlow}(q) = \sum_{i=1}^n \varphi_i RP(M_i) = \varphi \mathbf{R} \mathbf{1}^T < 0,$$

stating that the steady-state *mean potential fluid flow rate for the continuous place* q is negative. Stable infinite buffer models usually converge, hence, the existing steady-state PDF is also *unique* in this case.

We introduce the following steady-state probability functions, obtained from the transient ones by taking the limit $\delta \rightarrow \infty$.

- $\varphi_i = \lim_{\delta \rightarrow \infty} \mathbf{P}(M(\delta) = M_i)$ is the *steady-state discrete marking probability*;
- $\ell_i = \lim_{\delta \rightarrow \infty} \mathbf{P}(X(\delta) = 0, M(\delta) = M_i)$ is the *steady-state buffer empty probability (probability mass at the lower boundary)*;
- $F_i(x) = \lim_{\delta \rightarrow \infty} \mathbf{P}(X(\delta) < x, M(\delta) = M_i)$ is the *steady-state fluid probability distribution function*;
- $f_i(x) = \frac{dF_i(x)}{dx} = \lim_{h \rightarrow 0} \frac{F_i(x+h) - F_i(x)}{h} = \lim_{\delta \rightarrow \infty} \lim_{h \rightarrow 0} \frac{\mathbf{P}(x < X(\delta) < x+h, M(\delta) = M_i)}{h}$ is the *steady-state fluid probability density function*.

Let $\varphi, \ell, F(x), f(x)$ be the row vectors with the elements $\varphi_i, \ell_i, F_i(x), f_i(x)$, respectively ($1 \leq i \leq n$).

By the total probability law for the stationary behaviour, we have

$$\ell + \int_{0+}^{\infty} f(x)dx = \varphi.$$

The ordinary differential equations describing the stationary behaviour are

$$\frac{dF(x)}{dx} \mathbf{R} = F(x) \mathbf{Q}, \quad x > 0;$$

$$\frac{df(x)}{dx} \mathbf{R} = f(x) \mathbf{Q}, \quad x > 0.$$

Note that we have $\frac{dF(x)}{dx} = f(x)$, $F(0) = \ell$, $F(\infty) = \varphi$.

The ordinary differential equation for the steady-state buffer empty probabilities (stationary lower boundary conditions) are

$$f(0) \mathbf{R} = \ell \mathbf{Q}.$$

The stationary lower boundary constraint is: if $\mathcal{R}_{ii} = RP(M_i) > 0$ then $F_i(0) = \ell_i = 0$ ($1 \leq i \leq n$).

The stationary normalizing condition is

$$\ell \mathbf{1}^T + \int_{0+}^{\infty} f(x) dx \mathbf{1}^T = 1,$$

where $\mathbf{1}$ is a row vector of n values 1.

The solutions of the equations for $F(x)$ and $f(x)$ in the form of *matrix exponent* are $F(x) = \ell e^{x\mathbf{QR}^{-1}}$ and $f(x) = \ell \mathbf{QR}^{-1} e^{x\mathbf{QR}^{-1}}$, respectively. Since the steady-state existence implies boundedness of the SFM associated with an LFSPN and we do not have a finite upper fluid level bound, the positive eigenvalues of \mathbf{QR}^{-1} must be excluded. Moreover, \mathbf{R}^{-1} does not exist if for some i ($1 \leq i \leq n$) we have $\mathcal{R}_{ii} = 0$. These difficulties are avoided in the alternative solution method for $F(x)$, called *spectral decomposition* [55, 39, 28, 33, 30], which we outline below.

Let us define the sets of *negative discrete markings* of N as $DRS^-(N) = \{M \in DRS(N) \mid RP(M) < 0\}$, *zero discrete markings* of N as $DRS^0(N) = \{M \in DRS(N) \mid RP(M) = 0\}$ and *positive discrete markings* of N as $DRS^+(N) = \{M \in DRS(N) \mid RP(M) > 0\}$. The spectral decomposition is $F(x) = \sum_{j=1}^m a_j e^{\gamma_j x} v_j$, where a_j are scalar coefficients, γ_j are the eigenvalues and $v_j = (v_{j1}, \dots, v_{jn})$ are the eigenvectors of \mathbf{QR}^{-1} . Thus, each v_j is the solution of the equation $v_j(\mathbf{QR}^{-1} - \gamma_j \mathbf{I}) = 0$, where \mathbf{I} is the identity matrix of the order n , hence, $v_j(\mathbf{Q} - \gamma_j \mathbf{R}) = 0$.

Since for each non-zero v_j we must have $|\mathbf{Q} - \gamma_j \mathbf{R}| = 0$, the number of solutions $\gamma_1, \dots, \gamma_m$ is that of non-zero elements among $\mathcal{R}_{ii} = RP(M_i)$ ($1 \leq i \leq n$), i.e. $m = |DRS^-(N)| + |DRS^+(N)|$. We have 1 zero eigenvalue, $|DRS^+(N)|$ eigenvalues with a negative real part and $|DRS^-(N)| - 1$ eigenvalues with a positive real part. Let us reorder all the eigenvalues according to the sign of their real part (first, with zero one; then with negative one; at last, with positive one). The boundedness of $F(x)$ requires $a_j = 0$ if $Re(\gamma_j) > 0$ ($1 \leq j \leq m$). For the zero eigenvalue $\gamma_1 = 0$ we have $a_1 e^{\gamma_1 x} v_1 = a_1 v_1$, and for the corresponding eigenvector it holds $v_1 \mathbf{Q} = 0$. Then $F(x) = a_1 v_1 + \sum_{k=2}^{|DRS^+(N)|+1} a_k e^{\gamma_k x} v_k$, where $Re(\gamma_k) < 0$ ($2 \leq k \leq |DRS^+(N)| + 1$). Remember that $\varphi = F(\infty) = a_1 v_1$, hence, $F(x) = \varphi + \sum_{k=2}^{|DRS^+(N)|+1} a_k e^{\gamma_k x} v_k$.

It remains to find $|DRS^+(N)|$ coefficients a_k corresponding to the eigenvalues γ_k ($2 \leq k \leq |DRS^+(N)| + 1$). Remember the stationary lower boundary constraint: if $\mathcal{R}_{ll} = RP(M_l) > 0$ then $F_l(0) = \ell_l = 0$. Then for each $M_l \in DRS^+(N)$ we have $F_l(0) = \varphi_l + \sum_{k=2}^{|DRS^+(N)|+1} a_k v_{kl} = 0$. We get a system of $|DRS^+(N)|$ independent linear equations with $|DRS^+(N)|$ unknowns, for which a unique solution exists.

Then, using $F(x)$, we can find $f(x) = \frac{dF(x)}{dx}$ and $\ell = F(0)$.

Let N be an LFSPN. In [53], we have presented a number of steady-state *hybrid (discrete-continuous) performance indices (measures)*, which can be calculated based on the steady-state fluid probability density function $f(x)$ for the SFM of N [13, 31, 32, 29, 28, 40]. Note that the hybrid performance indices that do not depend on the fluid level coincide with the corresponding discrete performance measures.

5. FLUID TRACE EQUIVALENCE

Trace equivalences are the least discriminating ones. In the trace semantics, the behavior of a system is associated with the set of all possible sequences of actions, i.e. the protocols of computations. Thus, the points of choice of an external observer between several extensions of a particular computation are not taken into account.

The formal definition of fluid trace equivalence resembles that of ordinary Markovian trace equivalence, proposed on transition-labeled CTMCs in [62], on sequential and concurrent Markovian process calculi SMPC and CMPC in [8, 6, 7, 9] and on Uniform Labeled Transition Systems (ULtraS) in [10, 11]. While defining fluid trace equivalence, we additionally have to take into account the fluid flow rates in the corresponding discrete markings of two compared LFSPNs. Hence, in order to construct fluid trace equivalence, we should determine how to calculate the cumulative execution probabilities of all the specific (selected) paths. A *path* in the discrete reachability graph of an LFSPN is a sequence of its discrete markings and transitions that is generated by some firing sequence in the LFSPN.

First, we should *multiply the transition firing probabilities* for all the transitions along the paths starting in the initial discrete marking of the LFSPN. The resulting product will be the *execution probability of the path*. Second, we must *sum the path execution probabilities* for all the selected paths corresponding to the same *sequence of actions*, the same *sequence of the average sojourn times* and the same *sequence of the fluid flow rates* in all the discrete markings participating the paths. We suppose that each LFSPN has exactly one continuous place. The resulting sum will be the *cumulative execution probability of the selected paths* corresponding to some fluid stochastic trace. A *fluid stochastic trace* is a pair with the first element being the triple of the correlated sequences of actions, average sojourn times and fluid flow rates; the second element being the execution probability of the triple. Each element of the triple guarantees that fluid trace equivalence respects the following aspects of the LFSPNs behaviour: *functional activity*, *stochastic timing* and *fluid flow*.

Note that $CTMC(N)$ can be interpreted as a semi-Markov chain (SMC) [42], denoted by $SMC(N)$, which is analyzed by extracting from it the embedded (absorbing) discrete time Markov chain (EDTMC) corresponding to N , denoted by $EDTMC(N)$. The construction of the latter is analogous to that applied in the context of GSPNs in [44, 45, 4, 5]. $EDTMC(N)$ only describes the state changes of $SMC(N)$ while ignoring its time characteristics. Thus, to construct the EDTMC, we should abstract from all time aspects of behaviour of the SMC, i.e. from the sojourn time in its states. It is well-known that every SMC is fully described by the EDTMC and the state sojourn time distributions (the latter can be specified by the vector of PDFs of residence time in the states) [35].

We first propose some helpful definitions of the probability functions for the transition firings and discrete marking changes. Let N be an LFSPN, $M, \widetilde{M} \in DRS(N)$ be its discrete markings and $t \in Ena(M)$.

The (time-abstract) *probability that the transition t fires in M* is

$$PT(t, M) = \frac{\Omega_N(t, M)}{\sum_{u \in Ena(M)} \Omega_N(u, M)} = \frac{\Omega_N(t, M)}{RE(M)} = SJ(M)\Omega_N(t, M).$$

We have $\forall M \in \mathbb{N}^{|Pd_N|} \sum_{t \in Ena(M)} PT(t, M) = \sum_{t \in Ena(M)} \frac{\Omega_N(t, M)}{\sum_{u \in Ena(M)} \Omega_N(u, M)} = \frac{\sum_{t \in Ena(M)} \Omega_N(t, M)}{\sum_{u \in Ena(M)} \Omega_N(u, M)} = 1$, i.e. $PT(t, M)$ defines a probability distribution.

The *probability to move from M to \widetilde{M} by firing any transition* is

$$PM(M, \widetilde{M}) = \sum_{\{t|M \xrightarrow{t} \widetilde{M}\}} PT(t, M) = \frac{\sum_{\{t|M \xrightarrow{t} \widetilde{M}\}} \Omega_N(t)}{RE(M)} = SJ(M) \cdot \sum_{\{t|M \xrightarrow{t} \widetilde{M}\}} \Omega_N(t).$$

We write $M \rightarrow_{\mathcal{P}} \widetilde{M}$, if $M \rightarrow \widetilde{M}$, where $\mathcal{P} = PM(M, \widetilde{M})$. We have $\forall M \in \mathbb{N}^{|P^{d_N}|} \sum_{\{\widetilde{M}|M \rightarrow \widetilde{M}\}} PM(M, \widetilde{M}) = \sum_{\{\widetilde{M}|M \rightarrow \widetilde{M}\}} \sum_{\{t|M \xrightarrow{t} \widetilde{M}\}} PT(t, M) = \sum_{t \in E_{na}(M)} PT(t, M) = 1$, i.e. $PM(M, \widetilde{M})$ defines a probability distribution.

Definition 5. Let N be an LFSPN. The embedded (absorbing) discrete time Markov chain (EDTMC) of N , denoted by $EDTMC(N)$, has the state space $DRS(N)$, the initial state M_N and the transitions $M \rightarrow_{\mathcal{P}} \widetilde{M}$, if $M \rightarrow \widetilde{M}$, where $\mathcal{P} = PM(M, \widetilde{M})$.

The underlying SMC of N , denoted by $SMC(N)$, has the EDTMC $EDTMC(N)$ and the sojourn time in every $M \in DRS(N)$ is exponentially distributed with the parameter $RE(M)$.

Since the sojourn time in every $M \in DRS(N)$ is exponentially distributed, we have $SMC(N) = CTMC(N)$.

Let N be an LFSPN. The elements \mathcal{P}_{ij} ($1 \leq i, j \leq n = |DRS(N)|$) of the (one-step) transition probability matrix (TPM) \mathbf{P} for $EDTMC(N)$ are defined as

$$\mathcal{P}_{ij} = \begin{cases} PM(M_i, M_j), & M_i \rightarrow M_j; \\ 0, & \text{otherwise.} \end{cases}$$

Let X be a set, $n \in \mathbb{N}_{\geq 1}$ and $x_i \in X$ ($1 \leq i \leq n$). Then $\chi = x_1 \cdots x_n$ is a finite sequence over X of length $|\chi| = n$. When X is a set on numbers, we usually write $\chi = x_1 \circ \cdots \circ x_n$, to avoid confusion because of mixing up the operations of concatenation of sequences (\circ) and multiplication of numbers (\cdot). The empty sequence ε of length $|\varepsilon| = 0$ is an extra case. Let X^* denote the set of all finite sequences (including the empty one) over X .

Let $M_N = M_0 \xrightarrow{t_1} M_1 \xrightarrow{t_2} \cdots \xrightarrow{t_n} M_n$ ($n \in \mathbb{N}$) be a finite sequence of transition firings starting in the initial discrete marking M_N and called firing sequence in N . The firing sequence generates the path $M_0 t_1 M_1 t_2 \cdots t_n M_n$ in the discrete reachability graph $DRG(N)$. Since the first discrete marking $M_N = M_0$ of the path is fixed, one can see that the (finite) transition sequence $\vartheta = t_1 \cdots t_n$ in N uniquely determines the discrete marking sequence $M_0 \cdots M_n$, ending with the last discrete marking M_n of the mentioned path in $DRG(N)$. Hence, to refer the paths, one can simply use the transition sequences extracted from them as shown above. The empty transition sequence ε refers to the path M_0 , consisting just of one discrete marking (which is the first and last one of the path in such a case).

Let N be an LFSPN. The set of all (finite) transition sequences in N is

$$TranSeq(N) = \{\vartheta \mid \vartheta = \varepsilon \text{ or } \vartheta = t_1 \cdots t_n, M_N = M_0 \xrightarrow{t_1} M_1 \xrightarrow{t_2} \cdots \xrightarrow{t_n} M_n\}.$$

Let $\vartheta = t_1 \cdots t_n \in TranSeq(N)$ and $M_N = M_0 \xrightarrow{t_1} M_1 \xrightarrow{t_2} \cdots \xrightarrow{t_n} M_n$. The probability to execute the transition sequence ϑ is

$$PT(\vartheta) = \prod_{i=1}^n PT(t_i, M_{i-1}).$$

For $\vartheta = \varepsilon$ we define $PT(\varepsilon) = 1$. Let us prove that $\forall n \in \mathbb{N}$
 $\sum_{\{\vartheta \in TranSeq(N) \mid |\vartheta|=n\}} PT(\vartheta) = 1$, i.e. $PT(\vartheta)$ defines a probability distribution.

Lemma 1. *Let N be an LFSPN. Then $\forall n \in \mathbb{N}$*

$$\sum_{\{\vartheta \in TranSeq(N) \mid |\vartheta|=n\}} PT(\vartheta) = 1.$$

Proof. We prove by induction on the transition sequences length n .

- $n = 0$

By definition, $\sum_{\{\vartheta \in TranSeq(N) \mid |\vartheta|=0\}} PT(\vartheta) = PT(\varepsilon) = 1$.

- $n \rightarrow n + 1$

By distributivity law for multiplication and addition, and since

$$\begin{aligned} \forall M \in \mathbb{N}^{P^d N} \quad \sum_{t \in E_{na}(M)} PT(t, M) = 1, \text{ we get } \sum_{\{\vartheta \in TranSeq(N) \mid |\vartheta|=n+1\}} PT(\vartheta) = \\ \sum_{\{t_1, \dots, t_n, t_{n+1} \mid M_N = M_0 \xrightarrow{t_1} M_1 \xrightarrow{t_2} \dots \xrightarrow{t_n} M_n \xrightarrow{t_{n+1}} M_{n+1}\}} \prod_{i=1}^{n+1} PT(t_i, M_{i-1}) = \\ \sum_{\{t_1, \dots, t_n \mid M_N = M_0 \xrightarrow{t_1} M_1 \xrightarrow{t_2} \dots \xrightarrow{t_n} M_n\}} \sum_{\{t_{n+1} \mid M_n \xrightarrow{t_{n+1}} M_{n+1}\}} \prod_{i=1}^n PT(t_i, M_{i-1}) PT(t_{n+1}, M_n) = \\ \sum_{\{t_1, \dots, t_n \mid M_N = M_0 \xrightarrow{t_1} M_1 \xrightarrow{t_2} \dots \xrightarrow{t_n} M_n\}} \left(\prod_{i=1}^n PT(t_i, M_{i-1}) \sum_{\{t_{n+1} \mid M_n \xrightarrow{t_{n+1}} M_{n+1}\}} PT(t_{n+1}, M_n) \right) = \\ \sum_{\{t_1, \dots, t_n \mid M_N = M_0 \xrightarrow{t_1} M_1 \xrightarrow{t_2} \dots \xrightarrow{t_n} M_n\}} \prod_{i=1}^n PT(t_i, M_{i-1}) \cdot 1 = 1. \quad \square \end{aligned}$$

Let $\vartheta = t_1 \cdots t_n \in TranSeq(N)$ be a transition sequence in N and $M_N = M_0 \xrightarrow{t_1} M_1 \xrightarrow{t_2} \dots \xrightarrow{t_n} M_n$. The *action sequence* of ϑ is $L_N(\vartheta) = a_1 \cdots a_n \in Act^*$, where $L_N(t_i) = a_i$ ($1 \leq i \leq n$), i.e. it is the sequence of actions which label the transitions of that transition sequence. For $\vartheta = \varepsilon$ we define $L_N(\varepsilon) = \varepsilon$. Further, the *average sojourn time sequence* of $\vartheta = t_1 \cdots t_n$ is $SJ(\vartheta) = SJ(M_0) \circ \dots \circ SJ(M_n) \in \mathbb{R}_{>0}^*$, i.e. it is the sequence of average sojourn times in the discrete markings of the path to which ϑ refers. For $\vartheta = \varepsilon$ we define $SJ(\varepsilon) = SJ(M_0)$. Similarly, the *(potential) fluid flow rate sequence* of $\vartheta = t_1 \cdots t_n$ is $RP(\vartheta) = RP(M_0) \circ \dots \circ RP(M_n) \in \mathbb{R}^*$, i.e. it is the sequence of (potential) fluid flow rates in the discrete markings of the path to which ϑ refers. For $\vartheta = \varepsilon$ we define $RP(\varepsilon) = RP(M_0)$.

Let N be an LFSPN and $(\sigma, \varsigma, \varrho) \in Act^* \times \mathbb{R}_{>0}^* \times \mathbb{R}^*$. The set of $(\sigma, \varsigma, \varrho)$ -selected (finite) transition sequences in N is defined as

$$TranSeq(N, \sigma, \varsigma, \varrho) = \{\vartheta \in TranSeq(N) \mid L_N(\vartheta) = \sigma, SJ(\vartheta) = \varsigma, RP(\vartheta) = \varrho\}.$$

Let $TranSeq(N, \sigma, \varsigma, \varrho) \neq \emptyset$. Then the triple $(\sigma, \varsigma, \varrho)$, together with its execution probability, which is the cumulative execution probability of all the paths from which the triple is extracted (as described above), constitute a *fluid stochastic trace* of the LFSPN N . Fluid stochastic traces are formally introduced below, followed by the (first) definition of fluid stochastic trace equivalence.

Definition 6. *A (finite) fluid stochastic trace of an LFSPN N is a pair $((\sigma, \varsigma, \varrho), PT(\sigma, \varsigma, \varrho))$, where $TranSeq(N, \sigma, \varsigma, \varrho) \neq \emptyset$ and the (cumulative) probability to execute $(\sigma, \varsigma, \varrho)$ -selected transition sequences is*

$$PT(\sigma, \varsigma, \varrho) = \sum_{\vartheta \in TranSeq(N, \sigma, \varsigma, \varrho)} PT(\vartheta).$$

We denote the set of all fluid stochastic traces of an LFSPN N by $FluStochTraces(N)$. Two LFSPNs N and N' are fluid trace equivalent, denoted by $N \equiv_{fl} N'$, if

$$FluStochTraces(N) = FluStochTraces(N').$$

$PT(\sigma, \varsigma, \varrho)$ defines a probability distribution, since by Lemma 1, we have $\forall n \in \mathbb{N} \sum_{\{(\sigma, \varsigma, \varrho) \mid |\sigma|=n\}} PT(\sigma, \varsigma, \varrho) = \sum_{\{(\sigma, \varsigma, \varrho) \mid |\sigma|=n\}} \sum_{\vartheta \in TranSeq(N, \sigma, \varsigma, \varrho)} PT(\vartheta) = \sum_{(\sigma, \varsigma, \varrho)} \sum_{\{\vartheta \in TranSeq(N, \sigma, \varsigma, \varrho) \mid |\vartheta|=n\}} PT(\vartheta) = \sum_{\{\vartheta \in TranSeq(N) \mid |\vartheta|=n\}} PT(\vartheta) = 1$.

The following (second) definition of fluid stochastic trace equivalence does not use fluid stochastic traces.

Definition 7. Two LFSPNs N and N' are fluid trace equivalent, denoted by $N \equiv_{fl} N'$, if $\forall (\sigma, \varsigma, \varrho) \in Act^* \times \mathbb{R}_{>0}^* \times \mathbb{R}^*$ we have

$$\sum_{\vartheta \in TranSeq(N, \sigma, \varsigma, \varrho)} PT(\vartheta) = \sum_{\vartheta' \in TranSeq(N', \sigma, \varsigma, \varrho)} PT(\vartheta').$$

Note that in Definition 7, for $\vartheta = t_1 \cdots t_n \in TranSeq(N, \sigma, \varsigma, \varrho)$ with $M_N = M_0 \xrightarrow{t_1} M_1 \xrightarrow{t_2} \cdots \xrightarrow{t_n} M_n$ and $\vartheta' = t'_1 \cdots t'_n \in TranSeq(N', \sigma, \varsigma, \varrho)$ with $M_{N'} = M'_0 \xrightarrow{t'_1} M'_1 \xrightarrow{t'_2} \cdots \xrightarrow{t'_n} M'_n$, we have $PT(\vartheta) = \prod_{i=1}^n PT(t_i, M_{i-1}) = \prod_{i=1}^n SJ(M_{i-1})\Omega_N(t_i, M_{i-1})$ and $PT(\vartheta') = \prod_{i=1}^n PT(t'_i, M'_{i-1}) = \prod_{i=1}^n SJ(M'_{i-1})\Omega_N(t'_i, M'_{i-1})$. Then the equality $SJ(M_0) \circ \cdots \circ SJ(M_n) = SJ(\vartheta) = \varsigma = SJ(\vartheta') = SJ(M'_0) \circ \cdots \circ SJ(M'_n)$ implies that $\prod_{i=1}^n SJ(M_{i-1}) = \prod_{i=1}^n SJ(M'_{i-1})$. Hence, $PT(\vartheta) = PT(\vartheta')$ iff $\prod_{i=1}^n \Omega_N(t_i, M_{i-1}) = \prod_{i=1}^n \Omega_N(t'_i, M'_{i-1})$. This alternative equality results in the following (third) definition of fluid trace equivalence.

Definition 8. Two LFSPNs N and N' are fluid trace equivalent, denoted by $N \equiv_{fl} N'$, if $\forall (\sigma, \varsigma, \varrho) \in Act^* \times \mathbb{R}_{>0}^* \times \mathbb{R}^*$ we have

$$\sum_{\{t_1 \cdots t_n \in TranSeq(N, \sigma, \varsigma, \varrho) \mid M_N = M_0 \xrightarrow{t_1} M_1 \xrightarrow{t_2} \cdots \xrightarrow{t_n} M_n\}} \prod_{i=1}^n \Omega_N(t_i, M_{i-1}) = \sum_{\{t'_1 \cdots t'_n \in TranSeq(N', \sigma, \varsigma, \varrho) \mid M_{N'} = M'_0 \xrightarrow{t'_1} M'_1 \xrightarrow{t'_2} \cdots \xrightarrow{t'_n} M'_n\}} \prod_{i=1}^n \Omega_N(t'_i, M'_{i-1}).$$

Note that in the definition of $TranSeq(N, \sigma, \varsigma, \varrho)$, as well as in Definitions 6, 7 and 8, for $\vartheta \in T_N^*$, we may use the *exit rate sequences* $RE(\vartheta) = RE(M_0) \circ \cdots \circ RE(M_n) \in \mathbb{R}_{>0}^*$ instead of average sojourn time sequences $\varsigma = SJ(\vartheta) = SJ(M_0) \circ \cdots \circ SJ(M_n) \in \mathbb{R}_{>0}^*$, since we have $\forall M \in DRS(N) SJ(M) = \frac{1}{RE(M)}$ and $\forall M \in DRS(N) \forall M' \in DRS(N') SJ(M) = SJ(M') \Leftrightarrow RE(M) = RE(M')$.

Let N and N' be LFSPNs such that $Pc_N = \{q\}$ and $Pc_{N'} = \{q'\}$. Then the continuous place q' of N corresponds to q of N' , i.e. q and q' are the *respective* continuous places. For $M \in DRS(N)$ (or for $M' \in DRS(N')$) we denote by $RP(M)$ (or by $RP(M')$) the fluid level change rate for the continuous place q (or for the corresponding one q'), i.e. the argument discrete marking determines for which of the two continuous places, q or q' , the flow rate function RP is taken.

Let N be an LFSPN. The *average potential fluid change volume* in a continuous place $q \in Pc_N$ in the discrete marking $M \in DRS(N)$ is

$$FluidChange(q, M) = SJ(M)RP(M).$$

In order to define the probability function $PT(\sigma, \varsigma, \varrho)$, the transition sequences corresponding to a particular action sequence are also selected according to the specific average sojourn times and fluid flow rates in the discrete markings of the paths to which those transition sequences refer. One of several intuitions behind such an additional selection is as follows. The average potential fluid change volume in a continuous place q in the discrete marking M is a product of the average sojourn time and the constant (possibly zero or negative) potential fluid flow rate in M . In each of the corresponding discrete markings M and M' of the paths to which the corresponding transition sequences $\vartheta \in TranSeq(N, \sigma, \varsigma, \varrho)$ and $\vartheta' \in TranSeq(N', \sigma, \varsigma, \varrho)$ refer, we shall have the same average potential fluid change volume in the respective continuous places q and q' , i.e. $FluidChange(q, M) = SJ(M)RP(M) = SJ(M')RP(M') = FluidChange(q', M')$. Note that the average *actual* and *potential* fluid change volumes coincide unless the lower boundary of fluid in some continuous place is reached, setting hereupon the actual fluid flow rate in it equal to zero till the end of the sojourn time in the current discrete marking.

Note that our notion of fluid trace equivalence is based rather on that of Markovian trace equivalence from [62], since there the average sojourn times in the states “surrounding” the actions of the corresponding traces of the equivalent processes should *coincide* while in the definition of the mentioned equivalence from [8, 6, 7, 9], the shorter average sojourn time may simulate the longer one. If we would adopt such a simulation then the smaller fluid change volumes would model the bigger ones, since the potential fluid flow rate remains constant while residing in a discrete marking. Since we observe no intuition behind that modeling, we do not use it.

Let $\vartheta = t_1 \cdots t_n \in TranSeq(N)$ and $M_N = M_0 \xrightarrow{t_1} M_1 \xrightarrow{t_2} \cdots \xrightarrow{t_n} M_n$. The *average potential fluid change volume for the transition sequence* ϑ in a continuous place $q \in Pc_N$ is

$$FluidChange(q, \vartheta) = \sum_{i=0}^n FluidChange(q, M_i).$$

In [10, 11], the following two types of Markovian trace equivalence have been proposed. The *state-to-state* Markovian trace equivalence requires coincidence of average sojourn times in all corresponding discrete markings of the paths. The *end-to-end* Markovian trace equivalence demands that only the sums of average sojourn times for all corresponding discrete markings of the paths should be equal. As a basis for constructing fluid trace equivalence, we have taken the state-to-state relation, since the constant potential fluid flow rate in the discrete markings may differ with their change (moreover, the actual fluid flow rate function may become discontinuous when the lower fluid boundary for a continuous place is reached in some discrete marking). Therefore, while summing the potential fluid flow rates for all discrete markings of a path, an important information is lost. The information is needed to calculate the average potential fluid change volume for a transition sequence that refers to the path. The mentioned value is a sum of the average potential fluid change volumes for all corresponding discrete markings of the path. It coincides for the corresponding transition sequences $\vartheta \in TranSeq(N, \sigma, \varsigma, \varrho)$ and $\vartheta' \in TranSeq(N', \sigma, \varsigma, \varrho)$, i.e. $FluidChange(q, \vartheta) = FluidChange(q', \vartheta')$ for the respective continuous places q and q' . Again, note that the average *actual* and *potential* fluid change volumes for a transition sequence may differ, due to

discontinuity of the actual fluid flow rate functions for some discrete markings of the path to which the transition sequence refers.

Let $TranSeq(N, \sigma, \varsigma, \varrho) \neq \emptyset$. The *average potential fluid change volume for the* $(\sigma, \varsigma, \varrho)$ -*selected (finite) transition sequences* in a continuous place $q \in Pc_N$ is

$$FluidChange(q, (\sigma, \varsigma, \varrho)) = FluidChange(q, \vartheta) \forall \vartheta \in TranSeq(N, \sigma, \varsigma, \varrho).$$

Then, as mentioned above, for the respective continuous places q and q' of the LFSPNs N and N' , such that $TranSeq(N, \sigma, \varsigma, \varrho) \neq \emptyset \neq TranSeq(N', \sigma, \varsigma, \varrho)$, we have $FluidChange(q, (\sigma, \varsigma, \varrho)) = FluidChange(q', (\sigma, \varsigma, \varrho))$.

Let $n \in \mathbb{N}$. The *average potential fluid change volume for the transition sequences of length n* in a continuous place $q \in Pc_N$ is

$$FluidChange(q, n) = \sum_{\{\vartheta \in TranSeq(N) \mid |\vartheta|=n\}} FluidChange(q, \vartheta) PT(\vartheta).$$

We get $FluidChange(q, n) = \sum_{\{\vartheta \in TranSeq(N) \mid |\vartheta|=n\}} FluidChange(q, \vartheta) PT(\vartheta) = \sum_{\{(\sigma, \varsigma, \varrho) \mid TranSeq(N, \sigma, \varsigma, \varrho) \neq \emptyset \wedge |\sigma|=n\}} FluidChange(q, (\sigma, \varsigma, \varrho)) PT(\sigma, \varsigma, \varrho)$. For the respective continuous places q and q' of the LFSPNs N and N' with $N \equiv_{fl} N'$, we have $\forall n \in \mathbb{N} FluidChange(q, n) = FluidChange(q', n)$. Thus, fluid trace equivalence preserves average potential fluid change volume for the transition sequences of every certain length in the respective continuous places.

Example 1. In Figure 1, the LFSPNs N and N' are presented, such that $N \equiv_{fl} N'$. We have $DRS(N) = \{M_1, M_2\}$, where $M_1 = (1, 0)$, $M_2 = (0, 1)$, and $DRS(N') = \{M'_1, M'_2, M'_3\}$, where $M'_1 = (1, 0, 0)$, $M'_2 = (0, 1, 0)$, $M'_3 = (0, 0, 1)$.

In Figure 2, the discrete reachability graphs $DRG(N)$ and $DRG(N')$ are depicted. In Figure 3, the underlying CTMCs $CTMC(N)$ and $CTMC(N')$ are drawn. In Figure 4, the EDTMCs $EDTMC(N)$ and $EDTMC(N')$ are presented.

The sojourn time average and variance vectors of N are

$$SJ = \left(\frac{1}{2}, \frac{1}{2} \right), \quad VAR = \left(\frac{1}{4}, \frac{1}{4} \right).$$

The TRM \mathbf{Q} for $CTMC(N)$, TPM \mathbf{P} for $EDTMC(N)$ and FRM \mathbf{R} for the SFM of N are

$$\mathbf{Q} = \begin{pmatrix} -2 & 2 \\ 2 & -2 \end{pmatrix}, \quad \mathbf{P} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \mathbf{R} = \begin{pmatrix} 1 & 0 \\ 0 & -2 \end{pmatrix}.$$

The sojourn time average and variance vectors of N' are

$$SJ' = \left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2} \right), \quad VAR' = \left(\frac{1}{4}, \frac{1}{4}, \frac{1}{4} \right).$$

The TRM \mathbf{Q}' for $CTMC(N')$, TPM \mathbf{P}' for $EDTMC(N')$ and FRM \mathbf{R}' for the SFM of N' are

$$\mathbf{Q}' = \begin{pmatrix} -2 & 1 & 1 \\ 2 & -2 & 0 \\ 2 & 0 & -2 \end{pmatrix}, \quad \mathbf{P}' = \begin{pmatrix} 0 & \frac{1}{2} & \frac{1}{2} \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \quad \mathbf{R}' = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -2 & 0 \\ 0 & 0 & -2 \end{pmatrix}.$$




FIG. 1. Fluid trace equivalent LFSPNs

We have $t_1 t_2 \in \text{TranSeq}(N, ab, \frac{1}{2} \circ \frac{1}{2} \circ \frac{1}{2}, 1 \circ (-2) \circ 1)$ and $t_1 t_3 \in \text{TranSeq}(N, ac, \frac{1}{2} \circ \frac{1}{2} \circ \frac{1}{2}, 1 \circ (-2) \circ 1)$, hence, $\text{FluidChange}(q, t_1 t_2) = \text{FluidChange}(q, t_1 t_3) = \frac{1}{2} \cdot 1 + \frac{1}{2} \cdot (-2) + \frac{1}{2} \cdot 1 = 0$.

We have $t'_1 t'_3 \in \text{TranSeq}(N', ab, \frac{1}{2} \circ \frac{1}{2} \circ \frac{1}{2}, 1 \circ (-2) \circ 1)$ and $t'_2 t'_4 \in \text{TranSeq}(N', ac, \frac{1}{2} \circ \frac{1}{2} \circ \frac{1}{2}, 1 \circ (-2) \circ 1)$, hence, $\text{FluidChange}(q', t'_1 t'_3) = \text{FluidChange}(q', t'_2 t'_4) = \frac{1}{2} \cdot 1 + \frac{1}{2} \cdot (-2) + \frac{1}{2} \cdot 1 = 0$.

It holds $PT(t_1 t_2) = PT(t_1 t_3) = 1 \cdot \frac{1}{2} = \frac{1}{2}$ and $PT(t'_1 t'_3) = PT(t'_2 t'_4) = \frac{1}{2} \cdot 1 = \frac{1}{2}$.

We get $\text{FluStochTraces}(N) = \{((\varepsilon, \frac{1}{2}, 1), 1), ((a, \frac{1}{2} \circ \frac{1}{2}, 1 \circ (-2)), 1), ((ab, \frac{1}{2} \circ \frac{1}{2} \circ \frac{1}{2}, 1 \circ (-2) \circ 1), \frac{1}{2}), ((ac, \frac{1}{2} \circ \frac{1}{2} \circ \frac{1}{2}, 1 \circ (-2) \circ 1), \frac{1}{2}), \dots\} = \text{FluStochTraces}(N')$.

It holds $\text{FluidChange}(q, (a, \frac{1}{2} \circ \frac{1}{2}, 1 \circ (-2))) = \text{FluidChange}(q', (a, \frac{1}{2} \circ \frac{1}{2}, 1 \circ (-2))) = \frac{1}{2} \cdot 1 + \frac{1}{2} \cdot (-2) = -\frac{1}{2}$.

We then get $\text{FluidChange}(q, 1) = \text{FluidChange}(q, t_1)PT(t_1) = (-\frac{1}{2}) \cdot 1 = -\frac{1}{2} = (-\frac{1}{2}) \cdot \frac{1}{2} + (-\frac{1}{2}) \cdot \frac{1}{2} = \text{FluidChange}(q', t'_1)PT(t'_1) + \text{FluidChange}(q', t'_2)PT(t'_2) = \text{FluidChange}(q', 1)$.

In Figure 5, the ideal (since we have a stochastic process here, the actual and average sojourn times may differ) evolution of the actual fluid level for the continuous place q of the LFSPN N is depicted. One can see that $X(0.75) = 0$, i.e. at the time moment $\delta = 0.75$, the fluid level $X(\delta)$ reaches the zero low boundary while N resides in the discrete marking $M(\delta) = M_2$ for all $\delta \in [0.5; 1)$. Then the actual fluid flow rate function $RA(M(\delta), X(\delta))$ has a discontinuity at that point, where the function value is changed instantly from -2 to 0 . If it would exist no lower boundary, the average potential and actual fluid change volumes for the transition sequences of length 1 in the continuous place q would coincide and be equal to $\text{FluidChange}(q, 1) = -0.5 = 0.5 - 1 = X(1)$.

In Figure 6, possible evolution of the actual fluid level for the continuous place q of the LFSPN N is presented, where the actual and average sojourn times in the discrete markings demonstrate substantial differences.

6. FLUID BISIMULATION EQUIVALENCE

Bisimulation equivalences respect particular points of choice in the behavior of a system. To define fluid bisimulation equivalence, we have to consider a bisimulation




FIG. 2. The discrete reachability graphs of the fluid trace equivalent LFSPNs




FIG. 3. The underlying CTMCs of the fluid trace equivalent LFSPNs




FIG. 4. The EDTMCs of the fluid trace equivalent LFSPNs




FIG. 5. The ideal evolution of the actual fluid level in the first of two fluid trace equivalent LFSPNs




FIG. 6. Possible evolution of the actual fluid level in the first of two fluid trace equivalent LFSPNs

being an *equivalence* relation that partitions the states of the *union* of the discrete reachability graphs $DRG(N)$ and $DRG(N')$ of the LFSPNs N and N' . For N and N' to be bisimulation equivalent the initial states M_N and $M_{N'}$ of their discrete reachability graphs should be related by a bisimulation having the following transfer property: if two states are related then in each of them the same action can occur, leading with the identical overall rate from each of the two states to *the same equivalence class* for every such action.

The definition of fluid bisimulation should be given at the level of LFSPNs, but it must use the transition rates of the extracted CTMC. These rates cannot be easily (i.e. with a simple expression) defined at the level of more general LFSPNs, whose discrete part is labeled GSPNs. In addition, the action labels of immediate transitions are lost and their individual probabilities are redistributed while GSPNs are transformed into CTSPNs. The individual probabilities of immediate transitions are “dissolved” in the total transition rates between tangible states when vanishing states are eliminated from SMCs while reducing them to CTMCs. Therefore, to make the definition of fluid bisimulation less intricate, we have decided to consider only LFSPNs with labeled CTSPNs as the discrete part. Then the underlying stochastic process of the discrete part of LFSPNs will be that of CTSPNs, i.e. CTMCs.

The novelty of the fluid bisimulation definition w.r.t. that of the Markovian bisimulations from [15, 38, 8, 6, 7, 9, 10, 11] is that, for each pair of bisimilar discrete markings of N and N' , we require coincidence of the fluid flow rates of the *corresponding* (i.e. related by a *correspondence bijection*) continuous places of N and N' in these two discrete markings. Thus, fluid bisimulation equivalence takes into account *functional activity*, *stochastic timing* and *fluid flow*, like fluid trace equivalence does.

We first propose some helpful extensions of the rate functions for the discrete marking changes and for the fluid flow in continuous places. Let N be an LFSPN and $\mathcal{H} \subseteq DRS(N)$. Then, for each $M \in DRS(N)$ and $a \in Act$, we write $M \xrightarrow{a} \mathcal{H}$,

where $\lambda = RM_a(M, \mathcal{H})$ is the *overall rate to move from M into the set of discrete markings \mathcal{H} by action a* , defined as

$$RM_a(M, \mathcal{H}) = \sum_{\{t | \exists \widetilde{M} \in \mathcal{H} \ M \xrightarrow{t} \widetilde{M}, L_N(t)=a\}} \Omega_N(t, M).$$

We write $M \xrightarrow{a} \mathcal{H}$ if $\exists \lambda \ M \xrightarrow{a}_{\lambda} \mathcal{H}$. Further, we write $M \rightarrow_{\lambda} \mathcal{H}$ if $\exists a \ M \xrightarrow{a} \mathcal{H}$, where $\lambda = RM(M, \mathcal{H})$ is the *overall rate to move from M into the set of discrete markings \mathcal{H} by any actions*, defined as

$$RM(M, \mathcal{H}) = \sum_{\{t | \exists \widetilde{M} \in \mathcal{H} \ M \xrightarrow{t} \widetilde{M}\}} \Omega_N(t, M).$$

To construct a fluid bisimulation between LFSPNs N and N' , we should consider the “composite” set of their discrete markings $DRS(N) \cup DRS(N')$, since we have to identify the rates to come from any two equivalent discrete markings into the same “composite” equivalence class (w.r.t. the fluid bisimulation). Note that, for $N \neq N'$, transitions starting from the discrete markings of $DRS(N)$ (or $DRS(N')$) always lead to those from the same set, since $DRS(N) \cap DRS(N') = \emptyset$, and this allows us to “mix” the sets of discrete markings in the definition of fluid bisimulation.

Let $Pc_N = \{q\}$ and $Pc_{N'} = \{q'\}$. In this case the continuous place q' of N *corresponds* to q of N' , according to a trivial *correspondence bijection* $\beta : Pc_N \rightarrow Pc_{N'}$ such that $\beta(q) = q'$. Then for $M \in DRS(N)$ (or for $M' \in DRS(N')$) we denote by $RP(M)$ (or by $RP(M')$) the fluid level change rate for the continuous place q (or for the corresponding one q'), i.e. the argument discrete marking determines for which of the two continuous places, q or q' , the flow rate function RP is taken.

Note that if N and N' have more than one continuous place and there exists a *correspondence bijection* $\beta : Pc_N \rightarrow Pc_{N'}$ then we should consider several flow rate functions RP_i ($1 \leq i \leq l = |Pc_N| = |Pc_{N'}|$) in the same manner, i.e. each RP_i is used for the pair of the corresponding continuous places $q_i \in Pc_N$ and $\beta(q_i) = q'_i \in Pc_{N'}$. In other words, we require that the vectors $(RP_1(M), \dots, RP_l(M))$ and $(RP_1(M'), \dots, RP_l(M'))$ coincide for each pair of fluid bisimilar discrete markings M and M' in such a case.

Definition 9. *Let N and N' be LFSPNs such that $Pc_N = \{q\}$, $Pc_{N'} = \{q'\}$ and q' corresponds to q . An equivalence relation $\mathcal{R} \subseteq (DRS(N) \cup DRS(N'))^2$ is a fluid bisimulation between N and N' , denoted by $\mathcal{R} : N \xleftrightarrow{\text{fluid}} N'$, if:*

- (1) $(M_N, M_{N'}) \in \mathcal{R}$.
- (2) $(M_1, M_2) \in \mathcal{R} \Rightarrow RP(M_1) = RP(M_2), \forall \mathcal{H} \in (DRS(N) \cup DRS(N')) / \mathcal{R}, \forall a \in Act$

$$M_1 \xrightarrow{a}_{\lambda} \mathcal{H} \Leftrightarrow M_2 \xrightarrow{a}_{\lambda} \mathcal{H}.$$

Two LFSPNs N and N' are fluid bisimulation equivalent, denoted by $N \xleftrightarrow{\text{fluid}} N'$, if $\exists \mathcal{R} : N \xleftrightarrow{\text{fluid}} N'$.

Let $\mathcal{R}_{fl}(N, N') = \bigcup \{\mathcal{R} \mid \mathcal{R} : N \xleftrightarrow{\text{fluid}} N'\}$ be the *union of all fluid bisimulations* between N and N' . The following proposition proves that $\mathcal{R}_{fl}(N, N')$ is also an *equivalence* and $\mathcal{R}_{fl}(N, N') : N \xleftrightarrow{\text{fluid}} N'$.

Proposition 1. *Let N and N' be LFSPNs and $N \xleftrightarrow{\text{fluid}} N'$. Then $\mathcal{R}_{fl}(N, N')$ is the largest fluid bisimulation between N and N' .*

Proof. Analogous to that of Proposition 8.2.1 from [38], which establishes the result for strong equivalence. \square

Let N, N' be LFSPNs with $\mathcal{R} : N \leftrightarrow_{fl} N'$ and $\mathcal{H} \in (DRS(N) \cup DRS(N'))/\mathcal{R}$. We now present several remarks on the important equalities and helpful notations based on the rate functions RM_a , RM , RP and sojourn time characteristics SJ , VAR .

Remark 1. We have $\forall M_1, M_2 \in \mathcal{H} \forall \tilde{\mathcal{H}} \in (DRS(N) \cup DRS(N'))/\mathcal{R} \forall a \in Act M_1 \xrightarrow{a} \lambda \tilde{\mathcal{H}} \Leftrightarrow M_2 \xrightarrow{a} \lambda \tilde{\mathcal{H}}$. Since the previous equality is valid for all $M_1, M_2 \in \mathcal{H}$, we can rewrite it as $\mathcal{H} \xrightarrow{a} \lambda \tilde{\mathcal{H}}$, where $\lambda = RM_a(\mathcal{H}, \tilde{\mathcal{H}}) = RM_a(M_1, \tilde{\mathcal{H}}) = RM_a(M_2, \tilde{\mathcal{H}}) = RM_a(\mathcal{H} \cap DRS(N), \tilde{\mathcal{H}}) = RM_a(\mathcal{H} \cap DRS(N'), \tilde{\mathcal{H}})$. Then we write $\mathcal{H} \xrightarrow{a} \tilde{\mathcal{H}}$ if $\exists \lambda \mathcal{H} \xrightarrow{a} \lambda \tilde{\mathcal{H}}$ and $\mathcal{H} \rightarrow \tilde{\mathcal{H}}$ if $\exists a \mathcal{H} \xrightarrow{a} \tilde{\mathcal{H}}$.

Since the transitions from the discrete markings of $DRS(N)$ always lead to those from the same set, we have $\forall M \in DRS(N) \forall a \in Act RM_a(M, \tilde{\mathcal{H}}) = RM_a(M, \tilde{\mathcal{H}} \cap DRS(N))$. Hence, $\forall M \in \mathcal{H} \cap DRS(N) \forall a \in Act RM_a(\mathcal{H}, \tilde{\mathcal{H}}) = RM_a(M, \tilde{\mathcal{H}}) = RM_a(M, \tilde{\mathcal{H}} \cap DRS(N)) = RM_a(\mathcal{H} \cap DRS(N), \tilde{\mathcal{H}} \cap DRS(N))$. The same is true for $DRS(N')$. Thus, $\forall \tilde{\mathcal{H}} \in (DRS(N) \cup DRS(N'))/\mathcal{R}$

$$RM_a(\mathcal{H} \cap DRS(N), \tilde{\mathcal{H}} \cap DRS(N)) = RM_a(\mathcal{H}, \tilde{\mathcal{H}}) = RM_a(\mathcal{H} \cap DRS(N'), \tilde{\mathcal{H}} \cap DRS(N')).$$

Remark 2. We have $\forall M_1, M_2 \in \mathcal{H} \forall \tilde{\mathcal{H}} \in (DRS(N) \cup DRS(N'))/\mathcal{R} RM(M_1, \tilde{\mathcal{H}}) = \sum_{\{t | \exists \tilde{M}_1 \in \tilde{\mathcal{H}} M_1 \xrightarrow{t} \tilde{M}_1\}} \Omega_N(t, M_1) = \sum_{a \in Act} \sum_{\{t | \exists \tilde{M}_1 \in \tilde{\mathcal{H}} M_1 \xrightarrow{t} \tilde{M}_1, L_N(t)=a\}} \Omega_N(t, M_1) = \sum_{a \in Act} RM_a(M_1, \tilde{\mathcal{H}}) = \sum_{a \in Act} RM_a(M_2, \tilde{\mathcal{H}}) = \sum_{a \in Act} \sum_{\{t | \exists \tilde{M}_2 \in \tilde{\mathcal{H}} M_2 \xrightarrow{t} \tilde{M}_2, L_N(t)=a\}} \Omega_N(t, M_2) = \sum_{\{t | \exists \tilde{M}_2 \in \tilde{\mathcal{H}} M_2 \xrightarrow{t} \tilde{M}_2\}} \Omega_N(t, M_2) = RM(M_2, \tilde{\mathcal{H}})$. Since the previous equality is valid for all $M_1, M_2 \in \mathcal{H}$, we can denote $RM(\mathcal{H}, \tilde{\mathcal{H}}) = RM(M_1, \tilde{\mathcal{H}}) = RM(M_2, \tilde{\mathcal{H}})$. Then we write $\mathcal{H} \rightarrow \lambda \tilde{\mathcal{H}}$, where $\lambda = RM(\mathcal{H}, \tilde{\mathcal{H}}) = RM(M_1, \tilde{\mathcal{H}}) = RM(M_2, \tilde{\mathcal{H}})$.

Since the transitions from the discrete markings of $DRS(N)$ always lead to those from the same set, we have $\forall M \in DRS(N) RM(M, \tilde{\mathcal{H}}) = RM(M, \tilde{\mathcal{H}} \cap DRS(N))$. Hence, $\forall M \in \mathcal{H} \cap DRS(N) RM(\mathcal{H}, \tilde{\mathcal{H}}) = RM(M, \tilde{\mathcal{H}}) = RM(M, \tilde{\mathcal{H}} \cap DRS(N)) = RM(\mathcal{H} \cap DRS(N), \tilde{\mathcal{H}} \cap DRS(N))$. The same is true for $DRS(N')$. Thus, $\forall \tilde{\mathcal{H}} \in (DRS(N) \cup DRS(N'))/\mathcal{R}$

$$RM(\mathcal{H} \cap DRS(N), \tilde{\mathcal{H}} \cap DRS(N)) = RM(\mathcal{H}, \tilde{\mathcal{H}}) = RM(\mathcal{H} \cap DRS(N'), \tilde{\mathcal{H}} \cap DRS(N')).$$

Remark 3. We have $\forall M_1, M_2 \in \mathcal{H} RP(M_1) = RP(M_2)$. Since the previous equality is valid for all $M_1, M_2 \in \mathcal{H}$, we can denote $RP(\mathcal{H}) = RP(M_1) = RP(M_2)$.

Since any argument discrete marking $M \in DRS(N) \cup DRS(N')$ completely determines for which continuous place the flow rate function $RP(M)$ is taken (either for q if $M \in DRS(N)$ or for q' if $M \in DRS(N')$), we have $\forall M \in \mathcal{H} \cap DRS(N) RP(\mathcal{H}) = RP(M) = RP(\mathcal{H} \cap DRS(N))$. The same is true for $DRS(N')$. Thus,

$$RP(\mathcal{H} \cap DRS(N)) = RP(\mathcal{H}) = RP(\mathcal{H} \cap DRS(N')).$$

Remark 4. We have $\forall M_1, M_2 \in \mathcal{H} SJ(M_1) = \frac{1}{\sum_{t \in E_{na}(M_1)} \Omega_N(t, M_1)} = \frac{1}{\sum_{\tilde{\mathcal{H}} \in (DRS(N) \cup DRS(N'))/\mathcal{R}} \sum_{\{t | \exists \tilde{M}_1 \in \tilde{\mathcal{H}} M_1 \xrightarrow{t} \tilde{M}_1\}} \Omega_N(t, M_1)} = \frac{1}{\sum_{\tilde{\mathcal{H}} \in (DRS(N) \cup DRS(N'))/\mathcal{R}} RM(M_1, \tilde{\mathcal{H}})} = \frac{1}{\sum_{\tilde{\mathcal{H}} \in (DRS(N) \cup DRS(N'))/\mathcal{R}} RM(\mathcal{H}, \tilde{\mathcal{H}})}$

$$\frac{1}{\sum_{\tilde{\mathcal{H}} \in (DRS(N) \cup DRS(N'))/\mathcal{R}} RM(M_2, \tilde{\mathcal{H}})} = \frac{1}{\sum_{\tilde{\mathcal{H}} \in (DRS(N) \cup DRS(N'))/\mathcal{R}} \sum_{\{t | \exists \tilde{M}_2 \in \tilde{\mathcal{H}} \ M_2 \xrightarrow{t} \tilde{M}_2\}} \Omega_N(t, M_2)} = \frac{1}{\sum_{t \in E_{na}(M_2)} \Omega_N(t, M_2)} = SJ(M_2).$$

Since the previous equality is valid for all $M_1, M_2 \in \mathcal{H}$, we can denote $SJ_{\mathcal{R}}(\mathcal{H}) = SJ(M_1) = SJ(M_2)$.

Since any argument discrete marking $M \in DRS(N) \cup DRS(N')$ completely determines, for which LFSPN the average sojourn time function $SJ(M)$ is considered (for N if $M \in DRS(N)$, or for N' if $M \in DRS(N')$), we have $\forall M \in \mathcal{H} \cap DRS(N)$ $SJ(\mathcal{H}) = SJ(M) = SJ(\mathcal{H} \cap DRS(N))$. The same is true for $DRS(N')$. Thus,

$$SJ(\mathcal{H} \cap DRS(N)) = SJ(\mathcal{H}) = SJ(\mathcal{H} \cap DRS(N')).$$

Remark 5. We have $\forall M_1, M_2 \in \mathcal{H}$ $VAR(M_1) = \frac{1}{(\sum_{t \in E_{na}(M_1)} \Omega_N(t, M_1))^2} = \frac{1}{(\sum_{\tilde{\mathcal{H}} \in (DRS(N) \cup DRS(N'))/\mathcal{R}} \sum_{\{t | \exists \tilde{M}_1 \in \tilde{\mathcal{H}} \ M_1 \xrightarrow{t} \tilde{M}_1\}} \Omega_N(t, M_1))^2} = \frac{1}{(\sum_{\tilde{\mathcal{H}} \in (DRS(N) \cup DRS(N'))/\mathcal{R}} RM(M_1, \tilde{\mathcal{H}}))^2} = \frac{1}{(\sum_{\tilde{\mathcal{H}} \in (DRS(N) \cup DRS(N'))/\mathcal{R}} RM(\mathcal{H}, \tilde{\mathcal{H}}))^2} = \frac{1}{(\sum_{\tilde{\mathcal{H}} \in (DRS(N) \cup DRS(N'))/\mathcal{R}} RM(M_2, \tilde{\mathcal{H}}))^2} = \frac{1}{(\sum_{\tilde{\mathcal{H}} \in (DRS(N) \cup DRS(N'))/\mathcal{R}} \sum_{\{t | \exists \tilde{M}_2 \in \tilde{\mathcal{H}} \ M_2 \xrightarrow{t} \tilde{M}_2\}} \Omega_N(t, M_2))^2} = \frac{1}{(\sum_{t \in E_{na}(M_2)} \Omega_N(t, M_2))^2} = VAR(M_2)$. Since the previous equality is valid for all $M_1, M_2 \in \mathcal{H}$, we can denote $VAR_{\mathcal{R}}(\mathcal{H}) = VAR(M_1) = VAR(M_2)$.

Since any argument discrete marking $M \in DRS(N) \cup DRS(N')$ completely determines, for which LFSPN the sojourn time variance function $VAR(M)$ is considered (for N if $M \in DRS(N)$, or for N' if $M \in DRS(N')$), we have $\forall M \in \mathcal{H} \cap DRS(N)$ $VAR(\mathcal{H}) = VAR(M) = VAR(\mathcal{H} \cap DRS(N))$. The same is true for $DRS(N')$. Thus,

$$VAR(\mathcal{H} \cap DRS(N)) = VAR(\mathcal{H}) = VAR(\mathcal{H} \cap DRS(N')).$$

Example 2. In Figure 7, the LFSPNs N and N' are presented, such that $N \xleftrightarrow{fl} N'$. The only difference between the respective LFSPNs in Figure 1 and those in Figure 7 is that the transitions t_3 and t'_4 are labeled with action c in the former, instead of action b in the latter.

Therefore, the following notions coincide for the respective LFSPNs in Figure 1 and those in Figure 7: the discrete reachability sets $DRS(N)$ and $DRS(N')$, the discrete reachability graphs $DRG(N)$ and $DRG(N')$, the underlying CTMCs $CTMC(N)$ and $CTMC(N')$, the average sojourn time vectors SJ and SJ' of N and N' , the sojourn time variance vectors VAR and VAR' of N and N' , the TRMs \mathbf{Q} and \mathbf{Q}' for $CTMC(N)$ and $CTMC(N')$, the TPMs \mathbf{P} and \mathbf{P}' for $EDTMC(N)$ and $EDTMC(N')$, the FRMs \mathbf{R} and \mathbf{R}' for the SFMs of N and N' .

We have $(DRS(N) \cup DRS(N'))/\mathcal{R}_{fl(N, N')} = \{\mathcal{H}_1, \mathcal{H}_2\}$, where $\mathcal{H}_1 = \{M_1, M'_1\}$, $\mathcal{H}_2 = \{M_2, M'_2, M'_3\}$.

We now compare the fluid equivalences to discover their interrelations. The following proposition states that fluid bisimulation equivalence implies fluid trace one.

Proposition 2. For LFSPNs N and N' the following holds:

$$N \xleftrightarrow{fl} N' \Rightarrow N \equiv_{fl} N'.$$




FIG. 7. Fluid bisimulation equivalent LFSPNs

Proof. Let $\mathcal{R} : N \xleftrightarrow{fl} N'$, $\mathcal{H} \in (DRS(N) \cup DRS(N'))/\mathcal{R}$ and $M_1, M_2 \in \mathcal{H}$. We have $RP(M_1) = RP(M_2)$ and $\forall \tilde{\mathcal{H}} \in (DRS(N) \cup DRS(N'))/\mathcal{R} \forall a \in Act M_1 \xrightarrow{a} \lambda \tilde{\mathcal{H}} \Leftrightarrow M_2 \xrightarrow{a} \lambda \tilde{\mathcal{H}}$. Note that transitions from the discrete markings of $DRS(N)$ always lead to those from the same set, hence, $\forall M \in DRS(N) RM_a(M, \tilde{\mathcal{H}}) = RM_a(M, \tilde{\mathcal{H}} \cap DRS(N))$. The same is true for $DRS(N')$.

By Remark 1 from Section 6, we can write $\mathcal{H} \xrightarrow{a} \lambda \tilde{\mathcal{H}}$ and denote $\lambda = RM_a(M_1, \tilde{\mathcal{H}}) = RM_a(M_2, \tilde{\mathcal{H}}) = RM_a(\mathcal{H}, \tilde{\mathcal{H}}) = RM_a(\mathcal{H} \cap DRS(N), \tilde{\mathcal{H}} \cap DRS(N)) = RM_a(\mathcal{H} \cap DRS(N'), \tilde{\mathcal{H}} \cap DRS(N'))$.

Further, by Remark 4 from Section 6, we can denote $SJ(M_1) = SJ(M_2) = SJ(\mathcal{H}) = SJ(\mathcal{H} \cap DRS(N)) = SJ(\mathcal{H} \cap DRS(N'))$.

At last, by Remark 3 from Section 6, we can denote $RP(M_1) = RP(M_2) = RP(\mathcal{H}) = RP(\mathcal{H} \cap DRS(N)) = RP(\mathcal{H} \cap DRS(N'))$.

Let $TranSeq(N, \sigma, \varsigma, \varrho) \neq \emptyset$ and $\sigma = a_1 \cdots a_n \in Act^*$, $\varsigma = s_0 \circ \cdots \circ s_n \in \mathbb{R}_{>0}^*$, $\varrho = r_0 \circ \cdots \circ r_n \in \mathbb{R}^*$. Taking into account the notes above and $\mathcal{R} : N \xleftrightarrow{fl} N'$, we have $SJ(M_N) = SJ(M_{N'}) = s_0$, $RP(M_N) = RP(M_{N'}) = r_0$ and for all $\mathcal{H}_1, \dots, \mathcal{H}_n \in (DRS(N) \cup DRS(N'))/\mathcal{R}$, such that $SJ(\mathcal{H}_i) = s_i$, $RP(\mathcal{H}_i) = r_i$ ($1 \leq i \leq n$), it holds $M_N \xrightarrow{a_1} \lambda_1 \mathcal{H}_1 \xrightarrow{a_2} \lambda_2 \cdots \xrightarrow{a_n} \lambda_n \mathcal{H}_n \Leftrightarrow M_{N'} \xrightarrow{a_1} \lambda_1 \mathcal{H}_1 \xrightarrow{a_2} \lambda_2 \cdots \xrightarrow{a_n} \lambda_n \mathcal{H}_n$. Then we have $TranSeq(N', \sigma, \varsigma, \varrho) \neq \emptyset$. Thus, $TranSeq(N, \sigma, \varsigma, \varrho) \neq \emptyset$ implies $TranSeq(N', \sigma, \varsigma, \varrho) \neq \emptyset$.

We now intend to prove that the sum of the transition rates products for all the paths starting in $M_N = M_0$ and going through the discrete markings from $\mathcal{H}_1, \dots, \mathcal{H}_n$ is equal to the product of $\lambda_1, \dots, \lambda_n$, which is essentially the transition rates product for the “composite” path starting in $\mathcal{H}_0 = [M_0]_{\mathcal{R}}$ and going through the equivalence classes $\mathcal{H}_1, \dots, \mathcal{H}_n$ in $DRG(N)$:

$$\sum_{\{t_1, \dots, t_n | M_N \xrightarrow{t_1} \cdots \xrightarrow{t_n} M_n, L_N(t_i) = a_i, M_i \in \mathcal{H}_i (1 \leq i \leq n)\}} \prod_{i=1}^n \Omega_N(t_i, M_{i-1}) = \prod_{i=1}^n RM_{a_i}(\mathcal{H}_{i-1}, \mathcal{H}_i).$$

We prove this equality by induction on the “composite” path length n .

- $n = 1$

$$\sum_{\{t_1 | M_N = M_0 \xrightarrow{t_1} M_1, L_N(t_1) = a_1, M_1 \in \mathcal{H}_1\}} \Omega_N(t_1, M_0) = RM_{a_1}(M_0, \mathcal{H}_1) = RM_{a_1}(\mathcal{H}_0, \mathcal{H}_1).$$

- $n \rightarrow n + 1$

$$\equiv_{fl} \longleftarrow \xleftrightarrow{\quad} fl$$

FIG. 8. Interrelations of fluid equivalences

$$\begin{aligned}
& \sum_{\{t_1, \dots, t_n, t_{n+1} | M_N = M_0 \xrightarrow{t_1} \dots \xrightarrow{t_n} M_n \xrightarrow{t_{n+1}} M_{n+1}, L_N(t_i) = a_i, M_i \in \mathcal{H}_i \ (1 \leq i \leq n+1)\}} \prod_{i=1}^{n+1} \Omega_N(t_i, M_{i-1}) = \\
& \sum_{\{t_1, \dots, t_n | M_N = M_0 \xrightarrow{t_1} \dots \xrightarrow{t_n} M_n, L_N(t_i) = a_i, M_i \in \mathcal{H}_i \ (1 \leq i \leq n)\}} \prod_{i=1}^n \Omega_N(t_i, M_{i-1}) \Omega_N(t_{n+1}, M_n) = \\
& \sum_{\{t_1, \dots, t_n | M_N = M_0 \xrightarrow{t_1} \dots \xrightarrow{t_n} M_n, L_N(t_i) = a_i, M_i \in \mathcal{H}_i \ (1 \leq i \leq n)\}} \left[\prod_{i=1}^n \Omega_N(t_i, M_{i-1}) \sum_{\{t_{n+1} | M_n \xrightarrow{t_{n+1}} M_{n+1}, L_N(t_{n+1}) = a_{n+1}, M_n \in \mathcal{H}_n, M_{n+1} \in \mathcal{H}_{n+1}\}} \Omega_N(t_{n+1}, M_n) \right] = \\
& \sum_{\{t_1, \dots, t_n | M_N = M_0 \xrightarrow{t_1} \dots \xrightarrow{t_n} M_n, L_N(t_i) = a_i, M_i \in \mathcal{H}_i \ (1 \leq i \leq n)\}} \prod_{i=1}^n \Omega_N(t_i, M_{i-1}) RM_{a_{n+1}}(M_n, \mathcal{H}_{n+1}) = \\
& \sum_{\{t_1, \dots, t_n | M_N = M_0 \xrightarrow{t_1} \dots \xrightarrow{t_n} M_n, L_N(t_i) = a_i, M_i \in \mathcal{H}_i \ (1 \leq i \leq n)\}} \prod_{i=1}^n \Omega_N(t_i, M_{i-1}) RM_{a_{n+1}}(\mathcal{H}_n, \mathcal{H}_{n+1}) = \\
& RM_{a_{n+1}}(\mathcal{H}_n, \mathcal{H}_{n+1}) \sum_{\{t_1, \dots, t_n | M_N = M_0 \xrightarrow{t_1} \dots \xrightarrow{t_n} M_n, L_N(t_i) = a_i, M_i \in \mathcal{H}_i \ (1 \leq i \leq n)\}} \prod_{i=1}^n \Omega_N(t_i, M_{i-1}) = \\
& RM_{a_{n+1}}(\mathcal{H}_n, \mathcal{H}_{n+1}) \prod_{i=1}^n RM_{a_i}(\mathcal{H}_{i-1}, \mathcal{H}_i) = \prod_{i=1}^{n+1} RM_{a_i}(\mathcal{H}_{i-1}, \mathcal{H}_i).
\end{aligned}$$

Note that the equality that we have just proved can also be applied to N' .

One can see that the summation over *all* $(\sigma, \varsigma, \varrho)$ -selected transition sequences is the same as the summation over *all accordingly selected equivalence classes*:

$$\begin{aligned}
& \sum_{t_1 \dots t_n \in TranSeq(N, \sigma, \varsigma, \varrho)} \prod_{i=1}^n \Omega_N(t_i, M_{i-1}) = \\
& \sum_{\{t_1, \dots, t_n | M_N = M_0 \xrightarrow{t_1} \dots \xrightarrow{t_n} M_n, L_N(t_i) = a_i, SJ(M_i) = s_i, RP(M_j) = r_j \ (1 \leq i \leq n)\}} \prod_{i=1}^n \Omega_N(t_i, M_{i-1}) = \\
& \sum_{\{\mathcal{H}_1, \dots, \mathcal{H}_n | SJ(\mathcal{H}_i) = s_i, RP(\mathcal{H}_i) = r_i \ (1 \leq i \leq n)\}} \prod_{i=1}^n \Omega_N(t_i, M_{i-1}) = \\
& \sum_{\{t_1, \dots, t_n | M_N = M_0 \xrightarrow{t_1} \dots \xrightarrow{t_n} M_n, L_N(t_i) = a_i, M_i \in \mathcal{H}_i \ (1 \leq i \leq n)\}} \prod_{i=1}^n \Omega_N(t_i, M_{i-1}) = \\
& \sum_{\{\mathcal{H}_1, \dots, \mathcal{H}_n | SJ(\mathcal{H}_i) = s_i, RP(\mathcal{H}_i) = r_i \ (1 \leq i \leq n)\}} \prod_{i=1}^n RM_{a_i}(\mathcal{H}_{i-1}, \mathcal{H}_i) = \\
& \sum_{\{\mathcal{H}_1, \dots, \mathcal{H}_n | SJ(\mathcal{H}_i) = s_i, RP(\mathcal{H}_i) = r_i \ (1 \leq i \leq n)\}} \prod_{i=1}^n \Omega_{N'}(t'_i, M'_{i-1}) = \\
& \sum_{\{t'_1, \dots, t'_n | M_{N'} = M'_0 \xrightarrow{t'_1} \dots \xrightarrow{t'_n} M'_n, L_{N'}(t'_i) = a_i, M'_i \in \mathcal{H}_i \ (1 \leq i \leq n)\}} \prod_{i=1}^n \Omega_{N'}(t'_i, M'_{i-1}) = \\
& \sum_{\{t'_1, \dots, t'_n | M_{N'} = M'_0 \xrightarrow{t'_1} \dots \xrightarrow{t'_n} M'_n, L_{N'}(t'_i) = a_i, SJ(M'_i) = s_i, RP(M'_j) = r_j \ (1 \leq i \leq n)\}} \prod_{i=1}^n \Omega_{N'}(t'_i, M'_{i-1}) = \\
& \sum_{t'_1 \dots t'_n \in TranSeq(N', \sigma, \varsigma, \varrho)} \prod_{i=1}^n \Omega_{N'}(t'_i, M'_{i-1}). \text{ By the remark before Definition 8, the proba-} \\
& \text{bilities to execute } (\sigma, \varsigma, \varrho)\text{-selected transition sequences in } N \text{ and } N' \text{ coincide.}
\end{aligned}$$

We conclude that for all triples $(\sigma, \varsigma, \varrho) \in Act^* \times \mathbb{R}_{>0}^* \times \mathbb{R}^*$, it holds that $TranSeq(N, \sigma, \varsigma, \varrho) \neq \emptyset$ implies $TranSeq(N', \sigma, \varsigma, \varrho) \neq \emptyset$ and the execution probabilities of $(\sigma, \varsigma, \varrho)$ in N and N' are equal. The reverse implication is proved by symmetry of fluid bisimulation. \square

The following theorem compares discrimination power of the fluid equivalences.

Theorem 1. *For LFSPNs N and N' the following strict implication holds that is also depicted in Figure 8:*

$$N \xleftrightarrow{\quad}_{fl} N' \Rightarrow N \equiv_{fl} N'.$$

Proof. Let us check the validity of the implication.

- The implication $\xleftrightarrow{\quad}_{fl} \rightarrow \equiv_{fl}$ is valid by Proposition 2.

Let us see that the implication is strict, i.e. the reverse one does not work, by the following counterexample.

- In Figure 1, $N \equiv_{fl} N'$, but $N \not\equiv_{fl} N'$, since only in the LFSPN N' an action a can be executed so (by firing t'_2) that no action b can occur afterwards. \square

7. REDUCTION OF THE BEHAVIOUR

Fluid bisimulation equivalence can be used to reduce the discrete reachability graphs and underlying CTMCs of LFSPNs. Reductions of graph-based models, like transition systems (whose instances are reachability graphs and CTMCs), result in those with less states (the graph nodes). The goal of the reduction is to decrease the number of states in the semantic representation of the modeled system while preserving its important qualitative and quantitative properties. Thus, the reduction allows one to simplify the behavioural and performance analysis.

An *autobisimulation* is a bisimulation between an LFSPN and itself. Let N be an LFSPN with $\mathcal{R} : N \xleftrightarrow{fl} N$ and $\mathcal{K} \in DRS(N)/\mathcal{R}$.

Remarks 2, 4 and 5 from Section 6 allow us to present the following definitions. The *average sojourn time in the equivalence class (w.r.t. \mathcal{R}) of discrete markings \mathcal{K}* is

$$SJ_{\mathcal{R}}(\mathcal{K}) = \frac{1}{\sum_{\tilde{\mathcal{K}} \in DRS(N)/\mathcal{R}} RM(\mathcal{K}, \tilde{\mathcal{K}})} = SJ(M) \quad \forall M \in \mathcal{K}.$$

The *average sojourn time vector for the equivalence classes (w.r.t. \mathcal{R}) of discrete markings* of N , denoted by $SJ_{\mathcal{R}}$, has the elements $SJ_{\mathcal{R}}(\mathcal{K})$, $\mathcal{K} \in DRS(N)/\mathcal{R}$.

The *sojourn time variance in the equivalence class (w.r.t. \mathcal{R}) of discrete markings \mathcal{K}* is

$$VAR_{\mathcal{R}}(\mathcal{K}) = \frac{1}{\left(\sum_{\tilde{\mathcal{K}} \in DRS(N)/\mathcal{R}} RM(\mathcal{K}, \tilde{\mathcal{K}})\right)^2} = VAR(M) \quad \forall M \in \mathcal{K}.$$

The *sojourn time variance vector for the equivalence classes (w.r.t. \mathcal{R}) of discrete markings* of N , denoted by $VAR_{\mathcal{R}}$, has the elements $VAR_{\mathcal{R}}(\mathcal{K})$, $\mathcal{K} \in DRS(N)/\mathcal{R}$.

Let $\mathcal{R}_{fl}(N) = \bigcup \{\mathcal{R} \mid \mathcal{R} : N \xleftrightarrow{fl} N\}$ be the *union of all fluid autobisimulations* on N . By Proposition 1, $\mathcal{R}_{fl}(N)$ is the largest fluid autobisimulation on N . Based on the equivalence classes w.r.t. $\mathcal{R}_{fl}(N)$, the quotient (by \xleftrightarrow{fl}) discrete reachability graphs and quotient (by \xleftrightarrow{fl}) underlying CTMCs of LFSPNs can be defined. The mentioned equivalence classes become the quotient states. The average and variance for the sojourn time in a quotient state are those in the corresponding equivalence class, respectively. Every quotient transition between two such composite states represents all transitions (having the same action label in case of the discrete reachability graph quotient) from the first state to the second one.

Definition 10. *Let N be an LFSPN. The quotient (by \xleftrightarrow{fl}) discrete reachability graph of N is a labeled transition system $DRG_{\xleftrightarrow{fl}}(N) = (S_{\xleftrightarrow{fl}}, \mathcal{L}_{\xleftrightarrow{fl}}, \mathcal{T}_{\xleftrightarrow{fl}}, s_{\xleftrightarrow{fl}})$ with*

- $S_{\xleftrightarrow{fl}} = DRS(N)/\mathcal{R}_{fl}(N)$;
- $\mathcal{L}_{\xleftrightarrow{fl}} = Act \times \mathbb{R}_{>0}$;
- $\mathcal{T}_{\xleftrightarrow{fl}} = \{(\mathcal{K}, (a, RM_a(\mathcal{K}, \tilde{\mathcal{K}})), \tilde{\mathcal{K}}) \mid \mathcal{K}, \tilde{\mathcal{K}} \in DRS(N)/\mathcal{R}_{fl}(N), \mathcal{K} \xrightarrow{a} \tilde{\mathcal{K}}\}$;
- $s_{\xleftrightarrow{fl}} = [M_N]_{\mathcal{R}_{fl}(N)}$.

The transition $(\mathcal{K}, (a, \lambda), \tilde{\mathcal{K}}) \in \mathcal{T}_{\xleftrightarrow{fl}}$ will be written as $\mathcal{K} \xrightarrow{a, \lambda} \tilde{\mathcal{K}}$.

Let \simeq denote isomorphism between the quotient discrete reachability graphs that binds their initial states.

The quotient (by $\underline{\leftrightarrow}_{fl}$) average sojourn time vector of N is $SJ_{\underline{\leftrightarrow}_{fl}} = SJ_{\mathcal{R}_{fl}(N)}$. The quotient (by $\underline{\leftrightarrow}_{fl}$) sojourn time variance vector of N is $VAR_{\underline{\leftrightarrow}_{fl}} = VAR_{\mathcal{R}_{fl}(N)}$.

Definition 11. Let N be an LFSPN. The quotient (by $\underline{\leftrightarrow}_{fl}$) underlying CTMC of N , denoted by $CTMC_{\underline{\leftrightarrow}_{fl}}(N)$, has the state space $DRS(N)/\mathcal{R}_{fl}(N)$, the initial state $[M_N]_{\mathcal{R}_{fl}(N)}$ and the transitions $\mathcal{K} \rightarrow_\lambda \tilde{\mathcal{K}}$ if $\mathcal{K} \rightarrow \tilde{\mathcal{K}}$, where $\lambda = RM(\mathcal{K}, \tilde{\mathcal{K}})$.

The steady-state PMF $\varphi_{\underline{\leftrightarrow}_{fl}}$ for $CTMC_{\underline{\leftrightarrow}_{fl}}(N)$ is defined like the corresponding notion φ for $CTMC(N)$.

The quotients of both discrete reachability graphs and underlying CTMCs are the minimal reductions of the mentioned objects modulo fluid bisimulation. The quotients can be used to simplify analysis of system properties which are preserved by $\underline{\leftrightarrow}_{fl}$, since less states should be examined for it. Such a reduction method resembles that from [3] based on place bisimulation equivalence for Petri nets, excepting that the former method merges states, while the latter one merges places.

Let N be an LFSPN. We shall now construct the quotients (by $\underline{\leftrightarrow}_{fl}$) of the TRM for $CTMC(N)$, FRM for the associated SFM of N , average sojourn time vector and sojourn time variance vector of N , using special collector and distributor matrices. The quotient TRMs and FRMs are applied to describe the quotient associated SFMs of LFSPNs. Let $DRS(N) = \{M_1, \dots, M_n\}$ and $DRS(N)/\mathcal{R}_{fl}(N) = \{\mathcal{K}_1, \dots, \mathcal{K}_l\}$.

The elements $(\mathcal{Q}_{\underline{\leftrightarrow}_{fl}})_{rs}$ ($1 \leq r, s \leq l$) of the TRM $\mathbf{Q}_{\underline{\leftrightarrow}_{fl}}$ for $CTMC_{\underline{\leftrightarrow}_{fl}}(N)$ are

$$(\mathcal{Q}_{\underline{\leftrightarrow}_{fl}})_{rs} = \begin{cases} RM(\mathcal{K}_r, \mathcal{K}_s), & r \neq s; \\ -\sum_{\{k|1 \leq k \leq l, k \neq r\}} RM(\mathcal{K}_r, \mathcal{K}_k), & r = s. \end{cases}$$

Like it has been done for strong performance bisimulation on labeled CTSPNs in [15], the $l \times l$ TRM $\mathbf{Q}_{\underline{\leftrightarrow}_{fl}}$ for $CTMC_{\underline{\leftrightarrow}_{fl}}(N)$ can be constructed from the $n \times n$ TRM \mathbf{Q} for $CTMC(N)$ using the $n \times l$ collector matrix \mathbf{V} for the largest fluid autobisimulation $\mathcal{R}_{fl}(N)$ on N and the $l \times n$ distributor matrix \mathbf{W} for \mathbf{V} . Then \mathbf{W} should be a non-negative matrix (i.e. all its elements must be non-negative) with the elements of each its row summed to one, such that $\mathbf{WV} = \mathbf{I}$, where \mathbf{I} is the identity matrix of order l , i.e. \mathbf{W} is a *left-inverse matrix* for \mathbf{V} . It is known that for each collector matrix there is at least one distributor matrix, in particular, the matrix obtained by transposing \mathbf{V} and subsequent normalizing its rows, to guarantee that the elements of each row of the transposed matrix are summed to one.

The elements \mathcal{V}_{ir} ($1 \leq i \leq n$, $1 \leq r \leq l$) of the collector matrix \mathbf{V} for the largest fluid autobisimulation $\mathcal{R}_{fl}(N)$ on N are defined as

$$\mathcal{V}_{ir} = \begin{cases} 1, & M_i \in \mathcal{K}_r; \\ 0, & \text{otherwise.} \end{cases}$$

Thus, all the elements of \mathcal{V} are non-negative, as required. The row elements of \mathcal{V} are summed to one, since for each M_i ($1 \leq i \leq n$) there exists exactly one \mathcal{K}_r ($1 \leq r \leq l$) such that $M_i \in \mathcal{K}_r$. Hence,

$$\mathbf{V}\mathbf{1}^T = \mathbf{1}^T,$$

where $\mathbf{1}$ on the left side is the row vector of l values 1 while $\mathbf{1}$ on the right side is the row vector of n values 1.

For a vector $v = (v_1, \dots, v_l)$, let $Diag(v)$ be a diagonal matrix with the elements $Diag_{rs}(v)$ ($1 \leq r, s \leq l$) defined as

$$Diag_{rs}(v) = \begin{cases} v_r, & r = s; \\ 0, & \text{otherwise.} \end{cases}$$

The *distributor matrix* \mathbf{W} for the collector matrix \mathbf{V} is defined as

$$\mathbf{W} = (Diag(\mathbf{V}^T \mathbf{1}^T))^{-1} \mathbf{V}^T,$$

where $\mathbf{1}$ is the row vector of n values 1. One can check that $\mathbf{WV} = \mathbf{I}$, where \mathbf{I} is the identity matrix of order l .

The elements $(\mathcal{QV})_{is}$ ($1 \leq i \leq n$, $1 \leq s \leq l$) of the matrix \mathbf{QV} are

$$(\mathcal{QV})_{is} = \sum_{j=1}^n \mathcal{Q}_{ij} \mathcal{V}_{js} = \sum_{\{j | 1 \leq j \leq n, M_j \in \mathcal{K}_s\}} RM(M_i, M_j) = RM(M_i, \mathcal{K}_s).$$

As we know, for each M_i ($1 \leq i \leq n$) there exists exactly one \mathcal{K}_r ($1 \leq r \leq l$) such that $M_i \in \mathcal{K}_r$. By Remark 2 from Section 6, for all $M_i \in \mathcal{K}_r$ we have $RM(\mathcal{K}_r, \mathcal{K}_s) = RM(M_i, \mathcal{K}_s)$ ($1 \leq i \leq n$, $1 \leq r, s \leq l$). Then the elements $(\mathcal{VQ}_{\leftrightarrow fl})_{is}$ ($1 \leq i \leq n$, $1 \leq s \leq l$) of the matrix $\mathbf{VQ}_{\leftrightarrow fl}$ are

$$(\mathcal{VQ}_{\leftrightarrow fl})_{is} = \sum_{r=1}^l \mathcal{V}_{ir} (\mathcal{Q}_{\leftrightarrow fl})_{rs} = \sum_{\{r | 1 \leq r \leq l, M_i \in \mathcal{K}_r\}} RM(\mathcal{K}_r, \mathcal{K}_s) = RM(M_i, \mathcal{K}_s).$$

Therefore, we have

$$\mathbf{QV} = \mathbf{VQ}_{\leftrightarrow fl}, \quad \mathbf{WQV} = \mathbf{Q}_{\leftrightarrow fl}.$$

The elements $(\mathcal{R}_{\leftrightarrow fl})_{rs}$ ($1 \leq r, s \leq l$) of the FRM $\mathbf{R}_{\leftrightarrow fl}$ of the quotient (by $\leftrightarrow fl$) SFM of N for the continuous place q are defined as

$$(\mathcal{R}_{\leftrightarrow fl})_{rs} = \begin{cases} RP(\mathcal{K}_r), & r = s; \\ 0, & r \neq s. \end{cases}$$

Let \mathbf{R} be the FRM of the SFM of N for the continuous place q . The elements $(\mathcal{RV})_{is}$ ($1 \leq i \leq n$, $1 \leq s \leq l$) of the matrix \mathbf{RV} are

$$(\mathcal{RV})_{is} = \sum_{j=1}^n \mathcal{R}_{ij} \mathcal{V}_{js} = RP(M_i) \mathcal{V}_{is} = \begin{cases} RP(M_i), & M_i \in \mathcal{K}_s; \\ 0, & \text{otherwise.} \end{cases}$$

By Remark 2 from Section 6, for all $M_i \in \mathcal{K}_s$ we have $RP(\mathcal{K}_s) = RP(M_i)$ ($1 \leq i \leq n$, $1 \leq s \leq l$). Then the elements $(\mathcal{VR}_{\leftrightarrow fl})_{is}$ ($1 \leq i \leq n$, $1 \leq s \leq l$) of the matrix $\mathbf{VR}_{\leftrightarrow fl}$ are

$$(\mathcal{VR}_{\leftrightarrow fl})_{is} = \sum_{r=1}^l \mathcal{V}_{ir} (\mathcal{R}_{\leftrightarrow fl})_{rs} = \mathcal{V}_{is} RP(\mathcal{K}_s) = \begin{cases} RP(\mathcal{K}_s) = RP(M_i), & M_i \in \mathcal{K}_s; \\ 0, & \text{otherwise.} \end{cases}$$

Therefore, we also have

$$\mathbf{RV} = \mathbf{VR}_{\leftrightarrow fl}, \quad \mathbf{WRV} = \mathbf{R}_{\leftrightarrow fl}.$$

$$DRG_{\leftrightarrow_{fl}}(N) \quad DRG_{\leftrightarrow_{fl}}(N')$$




FIG. 9. The quotient discrete reachability graphs of the fluid bisimulation equivalent LFSPNs

Let us consider the matrices $Diag(SJ)$ and $Diag(SJ_{\leftrightarrow_{fl}})$. By analogy with the proven above for \mathbf{R} and $\mathbf{R}_{\leftrightarrow_{fl}}$, we can deduce $Diag(SJ)\mathbf{V} = \mathbf{V}Diag(SJ_{\leftrightarrow_{fl}})$ and $\mathbf{W}Diag(SJ)\mathbf{V} = Diag(SJ_{\leftrightarrow_{fl}})$. Therefore, we have

$$\mathbf{1W}Diag(SJ)\mathbf{V} = \mathbf{1}Diag(SJ_{\leftrightarrow_{fl}}) = SJ_{\leftrightarrow_{fl}},$$

where $\mathbf{1}$ is the row vector of l values 1. In a similar way, we obtain

$$\mathbf{1W}Diag(VAR)\mathbf{V} = \mathbf{1}Diag(VAR_{\leftrightarrow_{fl}}) = VAR_{\leftrightarrow_{fl}},$$

where $\mathbf{1}$ is the row vector of l values 1.

Example 3. Consider the LFSPNs N and N' in Figure 7, for which it holds $N_{\leftrightarrow_{fl}}N'$. We have $DRS(N)/\mathcal{R}_{fl}(N) = \{\mathcal{K}_1, \mathcal{K}_2\}$, where $\mathcal{K}_1 = \{M_1\}$, $\mathcal{K}_2 = \{M_2\}$, and $DRS(N')/\mathcal{R}_{fl}(N') = \{\mathcal{K}'_1, \mathcal{K}'_2\}$, where $\mathcal{K}'_1 = \{M'_1\}$, $\mathcal{K}'_2 = \{M'_2, M'_3\}$. In Figure 9, the quotient discrete reachability graphs $DRG_{\leftrightarrow_{fl}}(N)$ and $DRG_{\leftrightarrow_{fl}}(N')$ are depicted, for which we have $DRG_{\leftrightarrow_{fl}}(N) \simeq DRG_{\leftrightarrow_{fl}}(N')$. In Figure 10, the quotient underlying CTMCs $CTMC_{\leftrightarrow_{fl}}(N)$ and $CTMC_{\leftrightarrow_{fl}}(N')$ are drawn, for which it holds $CTMC_{\leftrightarrow_{fl}}(N) \simeq CTMC_{\leftrightarrow_{fl}}(N') \simeq CTMC(N)$. We have $\mathbf{Q}_{\leftrightarrow_{fl}} = \mathbf{Q}'_{\leftrightarrow_{fl}} = \mathbf{Q}$, $\mathbf{R}_{\leftrightarrow_{fl}} = \mathbf{R}'_{\leftrightarrow_{fl}} = \mathbf{R}$ and $SJ_{\leftrightarrow_{fl}} = SJ'_{\leftrightarrow_{fl}} = SJ$, $VAR_{\leftrightarrow_{fl}} = VAR'_{\leftrightarrow_{fl}} = VAR$.

The collector matrix \mathbf{V} for the largest fluid autobisimulation $\mathcal{R}_{fl}(N)$ on N and the distributor matrix \mathbf{W} for \mathbf{V} are

$$\mathbf{V} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 1 \end{pmatrix}, \quad \mathbf{W} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{2} & \frac{1}{2} \end{pmatrix}.$$

Then it is easy to check that

$$\mathbf{WQ}'\mathbf{V} = \mathbf{Q}, \quad \mathbf{WR}'\mathbf{V} = \mathbf{R}.$$

Hence, it holds that

$$\mathbf{1W}Diag(SJ')\mathbf{V} = SJ, \quad \mathbf{1W}Diag(VAR')\mathbf{V} = VAR,$$

$$\text{where } \mathbf{1} = (1, 1), \quad Diag(SJ') = \begin{pmatrix} \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 \\ 0 & 0 & \frac{1}{2} \end{pmatrix}, \quad Diag(VAR') = \begin{pmatrix} \frac{1}{4} & 0 & 0 \\ 0 & \frac{1}{4} & 0 \\ 0 & 0 & \frac{1}{4} \end{pmatrix}.$$

$$CTMC_{\underline{\leftrightarrow}_{fl}}(N) \quad CTMC_{\underline{\leftrightarrow}_{fl}}(N')$$




FIG. 10. The quotient underlying CTMCs of the fluid bisimulation equivalent LFSPNs

8. PRESERVATION OF THE QUANTITATIVE BEHAVIOUR

It is clear that the proposed fluid bisimulation equivalence of LFSPNs preserves their qualitative (functional) behaviour which is based on the actions assigned to the fired transitions. Let us examine if fluid bisimulation equivalence also preserves the quantitative (performance) behaviour of LFSPNs, taken for the steady states of their underlying CTMCs and associated SFMs. The quantitative behaviour takes into account the values of the rates and probabilities, as well as those of the related probability mass, distribution, density and mass at lower boundary functions. Then the quotients of the functions will describe the quotient (by $\underline{\leftrightarrow}_{fl}$) associated SFMs.

The next proposition shows that for two LFSPNs related by $\underline{\leftrightarrow}_{fl}$ their aggregate steady-state probabilities coincide for each equivalence class of discrete markings.

Proposition 3. *Let N, N' be LFSPNs with $\mathcal{R} : N \underline{\leftrightarrow}_{fl} N'$ and $\varphi = (\varphi_1, \dots, \varphi_n)$, $n = |DRS(N)|$, be the steady-state PMF for $CTMC(N)$ and $\varphi' = (\varphi'_1, \dots, \varphi'_m)$, $m = |DRS(N')|$, be the steady-state PMF for $CTMC(N')$. Then for all $\mathcal{H} \in (DRS(N) \cup DRS(N'))/\mathcal{R}$ we have*

$$\sum_{\{i|M_i \in \mathcal{H} \cap DRS(N)\}} \varphi_i = \sum_{\{j|M'_j \in \mathcal{H} \cap DRS(N')\}} \varphi'_j.$$

Proof. See [53]. □

Let N be an LFSPN and φ be the steady-state PMF for $CTMC(N)$. Let $\varphi_{\mathcal{K}}$, $\mathcal{K} \in DRS(N)/\mathcal{R}_{fl}(N)$, be the elements of the steady-state PMF for $CTMC_{\underline{\leftrightarrow}_{fl}}(N)$, denoted by $\varphi_{\underline{\leftrightarrow}_{fl}}$. By (the proof of) Proposition 3, for all $\mathcal{K} \in DRS(N)/\mathcal{R}_{fl}(N)$ we have

$$\varphi_{\mathcal{K}} = \sum_{\{i|M_i \in \mathcal{K}\}} \varphi_i.$$

Let \mathbf{V} be the collector matrix for the largest fluid autobisimulation $\mathcal{R}_{fl}(N)$ on N . One can see that

$$\varphi \mathbf{V} = \varphi_{\underline{\leftrightarrow}_{fl}}.$$

We have $\begin{cases} \varphi \mathbf{Q} = \mathbf{0} \\ \varphi \mathbf{1}^T = 1 \end{cases}$. After right-multiplying both sides of the first equation by \mathbf{V} and since $\mathbf{V} \mathbf{1}^T = \mathbf{1}^T$, we get $\begin{cases} \varphi \mathbf{Q} \mathbf{V} = \mathbf{0} \\ \varphi \mathbf{V} \mathbf{1}^T = 1 \end{cases}$. Since $\mathbf{Q} \mathbf{V} = \mathbf{V} \mathbf{Q}_{\leftrightarrow_{fl}}$, we obtain $\begin{cases} \varphi \mathbf{V} \mathbf{Q}_{\leftrightarrow_{fl}} = \mathbf{0} \\ \varphi \mathbf{V} \mathbf{1}^T = 1 \end{cases}$. Since $\varphi \mathbf{V} = \varphi_{\leftrightarrow_{fl}}$, $\varphi_{\leftrightarrow_{fl}}$ is a solution of the linear equation system
$$\begin{cases} \varphi_{\leftrightarrow_{fl}} \mathbf{Q}_{\leftrightarrow_{fl}} = \mathbf{0} \\ \varphi_{\leftrightarrow_{fl}} \mathbf{1}^T = 1 \end{cases}.$$

Thus, the treatment of $CTMC_{\leftrightarrow_{fl}}(N)$ instead of $CTMC(N)$ simplifies the analytical solution, since we have less states, but constructing the TRM $\mathbf{Q}_{\leftrightarrow_{fl}}$ for $CTMC_{\leftrightarrow_{fl}}(N)$ also requires some efforts, including determining $\mathcal{R}_{fl}(N)$ and calculating the rates to move from one equivalence class to another. The behaviour of $CTMC_{\leftrightarrow_{fl}}(N)$ stabilizes quicker than that of $CTMC(N)$ (if each of them has a single steady state), since $\mathbf{Q}_{\leftrightarrow_{fl}}$ is denser matrix than \mathbf{Q} (the TRM for $CTMC(N)$) due to the fact that the former matrix is smaller and the transitions between the equivalence classes “include” all the transitions between the discrete markings belonging to these equivalence classes.

The following proposition shows that for two LFSPNs related by \leftrightarrow_{fl} their aggregate steady-state fluid PDFs coincide for each equivalence class of discrete markings.

Proposition 4. *Let N, N' be LFSPNs with $\mathcal{R} : N \leftrightarrow_{fl} N'$ and $F(x) = (F_1(x), \dots, F_n(x))$, $n = |DRS(N)|$, be the steady-state fluid PDF for the SFM of N and $F'(x) = (F'_1(x), \dots, F'_m(x))$, $m = |DRS(N')|$, be the steady-state fluid PDF for the SFM of N' . Then for all $\mathcal{H} \in (DRS(N) \cup DRS(N'))/\mathcal{R}$ we have*

$$\sum_{\{i|M_i \in \mathcal{H} \cap DRS(N)\}} F_i(x) = \sum_{\{j|M'_j \in \mathcal{H} \cap DRS(N')\}} F'_j(x), \quad x > 0.$$

Proof. See [53]. □

Let N be an LFSPN and $F(x)$ be the steady-state fluid PDF for the SFM of N . Let $F_{\mathcal{K}}(x)$, $\mathcal{K} \in DRS(N)/\mathcal{R}_{fl}(N)$, be the elements of the steady-state fluid PDF for the quotient (by \leftrightarrow_{fl}) SFM of N , denoted by $F_{\leftrightarrow_{fl}}(x)$. By (the proof of) Proposition 4, for all $\mathcal{K} \in DRS(N)/\mathcal{R}_{fl}(N)$ we have

$$F_{\mathcal{K}}(x) = \sum_{\{i|M_i \in \mathcal{K}\}} F_i(x), \quad x > 0.$$

Let \mathbf{V} be the collector matrix for the largest fluid autobisimulation $\mathcal{R}_{fl}(N)$ on N . One can see that

$$F(x) \mathbf{V} = F_{\leftrightarrow_{fl}}(x), \quad x > 0.$$

We have $\frac{dF(x)}{dx} \mathbf{R} = F(x) \mathbf{Q}$, $x > 0$. After right-multiplying both sides of the above equation by \mathbf{V} , we get $\frac{dF(x)}{dx} \mathbf{R} \mathbf{V} = F(x) \mathbf{Q} \mathbf{V}$, $x > 0$. Since $\mathbf{R} \mathbf{V} = \mathbf{V} \mathbf{R}_{\leftrightarrow_{fl}}$ and $\mathbf{Q} \mathbf{V} = \mathbf{V} \mathbf{Q}_{\leftrightarrow_{fl}}$, we obtain $\frac{dF(x)}{dx} \mathbf{V} \mathbf{R}_{\leftrightarrow_{fl}} = F(x) \mathbf{V} \mathbf{Q}_{\leftrightarrow_{fl}}$, $x > 0$. By linearity of differentiation operator, we have $\frac{d}{dx}(F(x) \mathbf{V}) \mathbf{R}_{\leftrightarrow_{fl}} = F(x) \mathbf{V} \mathbf{Q}_{\leftrightarrow_{fl}}$, $x > 0$. Since $F(x) \mathbf{V} = F_{\leftrightarrow_{fl}}(x)$, we conclude that $F_{\leftrightarrow_{fl}}(x)$ is a solution of the system of ordinary differential equations

$$\frac{dF_{\leftrightarrow_{fl}}(x)}{dx} \mathbf{R}_{\leftrightarrow_{fl}} = F_{\leftrightarrow_{fl}}(x) \mathbf{Q}_{\leftrightarrow_{fl}}, \quad x > 0.$$

Thus, the treatment of the quotient (by \leftrightarrow_{fl}) SFM of N instead of SFM of N simplifies the analytical solution.

The following proposition demonstrates that for two LFSPNs related by \leftrightarrow_{fl} their aggregate steady-state fluid probability density functions coincide for each equivalence class of discrete markings.

Proposition 5. *Let N, N' be LFSPNs with $\mathcal{R} : N \leftrightarrow_{fl} N'$ and $f(x) = (f_1(x), \dots, f_n(x))$, $n = |DRS(N)|$, be the steady-state fluid probability density function for the SFM of N and $f'(x) = (f'_1(x), \dots, f'_m(x))$, $m = |DRS(N')|$, be the steady-state fluid probability density function for the SFM of N' . Then for all $\mathcal{H} \in (DRS(N) \cup DRS(N'))/\mathcal{R}$ we have*

$$\sum_{\{i|M_i \in \mathcal{H} \cap DRS(N)\}} f_i(x) = \sum_{\{j|M'_j \in \mathcal{H} \cap DRS(N')\}} f'_j(x), \quad x > 0.$$

Proof. See [53]. □

Let N be an LFSPN and $f(x)$ be the steady-state fluid probability density function for the SFM of N . Let $f_{\mathcal{K}}(x)$, $\mathcal{K} \in DRS(N)/\mathcal{R}_{fl}(N)$, be the elements of the steady-state fluid probability density function for the quotient (by \leftrightarrow_{fl}) SFM of N , denoted by $f_{\leftrightarrow_{fl}}(x)$. By (the proof of) Proposition 5, for all $\mathcal{K} \in DRS(N)/\mathcal{R}_{fl}(N)$ we have

$$f_{\mathcal{K}}(x) = \sum_{\{i|M_i \in \mathcal{K}\}} f_i(x), \quad x > 0.$$

Let \mathbf{V} be the collector matrix for the largest fluid autobisimulation $\mathcal{R}_{fl}(N)$ on N . One can see that

$$f(x) \mathbf{V} = f_{\leftrightarrow_{fl}}(x), \quad x > 0.$$

We have $\frac{df(x)}{dx} \mathbf{R} = f(x) \mathbf{Q}$, $x > 0$. Like it has been done after Proposition 4, we can prove that $f_{\leftrightarrow_{fl}}(x)$ is a solution of the system of ordinary differential equations

$$\frac{df_{\leftrightarrow_{fl}}(x)}{dx} \mathbf{R}_{\leftrightarrow_{fl}} = f_{\leftrightarrow_{fl}}(x) \mathbf{Q}_{\leftrightarrow_{fl}}, \quad x > 0.$$

The following proposition demonstrates that for two LFSPNs related by \leftrightarrow_{fl} their aggregate steady-state buffer empty probabilities coincide for each equivalence class of discrete markings.

Proposition 6. *Let N, N' be LFSPNs with $\mathcal{R} : N \leftrightarrow_{fl} N'$ and $\ell = (\ell_1, \dots, \ell_n)$, $n = |DRS(N)|$, be the steady-state buffer empty probability for the SFM of N and $\ell'(x) = (\ell'_1, \dots, \ell'_m)$, $m = |DRS(N')|$, be the steady-state buffer empty probability for the SFM of N' . Then for all $\mathcal{H} \in (DRS(N) \cup DRS(N'))/\mathcal{R}$ we have*

$$\sum_{\{i|M_i \in \mathcal{H} \cap DRS(N)\}} \ell_i = \sum_{\{j|M'_j \in \mathcal{H} \cap DRS(N')\}} \ell'_j.$$

Proof. See [53]. □

Let N be an LFSPN and ℓ be the steady-state buffer empty probability for the SFM of N . Let $\ell_{\mathcal{K}}$, $\mathcal{K} \in DRS(N)/\mathcal{R}_{fl}(N)$, be the elements of the steady-state buffer empty probability for the quotient (by $\underline{\leftrightarrow}_{fl}$) SFM of N , denoted by $\ell_{\underline{\leftrightarrow}_{fl}}$. By (the proof of) Proposition 6, for all $\mathcal{K} \in DRS(N)/\mathcal{R}_{fl}(N)$ we have

$$\ell_{\mathcal{K}} = \sum_{\{i|M_i \in \mathcal{K}\}} \ell_i.$$

Let \mathbf{V} be the collector matrix for the largest fluid autobisimulation $\mathcal{R}_{fl}(N)$ on N . One can see that

$$\ell \mathbf{V} = \ell_{\underline{\leftrightarrow}_{fl}}.$$

We have $\ell = \varphi - \int_{0+}^{\infty} f(x)dx$. After right-multiplying both sides of the equation by \mathbf{V} , we get $\ell \mathbf{V} = \varphi \mathbf{V} - \left(\int_{0+}^{\infty} f(x)dx \right) \mathbf{V}$. Since $\ell \mathbf{V} = \ell_{\underline{\leftrightarrow}_{fl}}$ and $\varphi \mathbf{V} = \varphi_{\underline{\leftrightarrow}_{fl}}$, by linearity of integration operator, we obtain $\ell_{\underline{\leftrightarrow}_{fl}} = \varphi_{\underline{\leftrightarrow}_{fl}} - \int_{0+}^{\infty} f(x) \mathbf{V} dx$. Since $f(x) \mathbf{V} = f_{\underline{\leftrightarrow}_{fl}}(x)$, $x > 0$, then $\ell_{\underline{\leftrightarrow}_{fl}}$ is a solution of the linear equation system

$$\ell_{\underline{\leftrightarrow}_{fl}} = \varphi_{\underline{\leftrightarrow}_{fl}} - \int_{0+}^{\infty} f_{\underline{\leftrightarrow}_{fl}}(x) dx.$$

Thus, the proposed quotients of the probability functions describe the behaviour of the quotient (by $\underline{\leftrightarrow}_{fl}$) associated SFMs of LFSPNs.

Example 4. Consider the LFSPNs N and N' in Figure 7, for which it holds $N \underline{\leftrightarrow}_{fl} N'$. We have $DRS^-(N) = \{M_2\}$, $DRS^0(N) = \emptyset$ and $DRS^+(N) = \{M_1\}$.

The steady-state PMF for CTMC(N) is

$$\varphi = \left(\frac{1}{2}, \frac{1}{2} \right).$$

Then the stability condition for the SFM of N is fulfilled: $FluidFlow(q) = \sum_{i=1}^2 \varphi_i RP(M_i) = \frac{1}{2} \cdot 1 + \frac{1}{2}(-2) = -\frac{1}{2} < 0$.

For each eigenvalue γ we must have $|\gamma \mathbf{R} - \mathbf{Q}| = \begin{vmatrix} \gamma + 2 & -2 \\ -2 & -2\gamma + 2 \end{vmatrix} = -2\gamma(1 + \gamma) = 0$; hence, $\gamma_1 = 0$ and $\gamma_2 = -1$.

The corresponding eigenvectors are the solutions of

$$v_1 \begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix} = 0, \quad v_2 \begin{pmatrix} 1 & -2 \\ -2 & 4 \end{pmatrix} = 0.$$

Then the eigenvectors are $v_1 = \left(\frac{1}{2}, \frac{1}{2} \right)$ and $v_2 = \left(\frac{2}{3}, \frac{1}{3} \right)$.

Since $\varphi = F(\infty) = a_1 v_1$, we have $F(x) = \varphi + a_2 e^{\gamma_2 x} v_2$ and $a_1 = 1$. Since $\forall M_i \in DRS^+(N) F_i(0) = \varphi_i + a_2 v_{2i} = 0$ and $DRS^+(N) = \{M_1\}$, we have $\varphi_1 + a_2 v_{21} = \frac{1}{2} + a_2 \frac{2}{3} = 0$; hence, $a_2 = -\frac{3}{4}$.

Then the steady-state fluid PDF for the SFM of N is

$$F(x) = \left(\frac{1}{2} - \frac{1}{2} e^{-x}, \frac{1}{2} - \frac{1}{4} e^{-x} \right).$$

The steady-state fluid probability density function for the SFM of N is

$$f(x) = \frac{dF(x)}{dx} = \left(\frac{1}{2} e^{-x}, \frac{1}{4} e^{-x} \right).$$

The steady-state buffer empty probability for the SFM of N is

$$\ell = F(0) = \left(0, \frac{1}{4}\right).$$

We have $DRS^-(N') = \{M'_2, M'_3\}$, $DRS^0(N') = \emptyset$ and $DRS^+(N') = \{M'_1\}$.
The steady-state PMF for CTMC(N') is

$$\varphi' = \left(\frac{1}{2}, \frac{1}{4}, \frac{1}{4}\right).$$

Then the stability condition for the SFM of N' is fulfilled: $FluidFlow(q') = \sum_{j=1}^3 \varphi'_j RP(M'_j) = \frac{1}{2} \cdot 1 + \frac{1}{4}(-2) + \frac{1}{4}(-2) = -\frac{1}{2} < 0$.

For each eigenvalue γ' we must have $|\gamma' \mathbf{R}' - \mathbf{Q}'| = \begin{vmatrix} \gamma' + 2 & -1 & -1 \\ -2 & -2\gamma' + 2 & 0 \\ -2 & 0 & -2\gamma' + 2 \end{vmatrix} =$

$-2\gamma'(1 + \gamma')(1 - \gamma') = 0$; hence, $\gamma'_1 = 0$, $\gamma'_2 = -1$ and $\gamma'_3 = 1$.

By the boundedness condition, the positive eigenvalue γ'_3 and the corresponding eigenvector v'_3 should be excluded from the solution.

The remaining corresponding eigenvectors are the solutions of

$$v'_1 \begin{pmatrix} 2 & -1 & -1 \\ -2 & 2 & 0 \\ -2 & 0 & 2 \end{pmatrix} = 0, \quad v'_2 \begin{pmatrix} 1 & -1 & -1 \\ -2 & 4 & 0 \\ -2 & 0 & 4 \end{pmatrix} = 0.$$

Then the remaining eigenvectors are $v'_1 = (\frac{1}{2}, \frac{1}{4}, \frac{1}{4})$ and $v'_2 = (\frac{2}{3}, \frac{1}{6}, \frac{1}{6})$.

Since $\varphi' = F'(\infty) = a'_1 v'_1$, we have $F'(x) = \varphi' + a'_2 e^{\gamma'_2 x} v'_2$ and $a'_1 = 1$. Since $\forall M'_i \in DRS^+(N')$ $F'_i(0) = \varphi'_i + a'_2 v'_{2i} = 0$ and $DRS^+(N') = \{M'_1\}$, we have $\varphi'_1 + a'_2 v'_{21} = \frac{1}{2} + a'_2 \frac{2}{3} = 0$; hence, $a_2 = -\frac{3}{4}$.

Then the steady-state fluid PDF for the SFM of N' is

$$F'(x) = \left(\frac{1}{2} - \frac{1}{2}e^{-x}, \frac{1}{4} - \frac{1}{8}e^{-x}, \frac{1}{4} - \frac{1}{8}e^{-x}\right).$$

The steady-state fluid probability density function for the SFM of N' is

$$f'(x) = \frac{dF'(x)}{dx} = \left(\frac{1}{2}e^{-x}, \frac{1}{8}e^{-x}, \frac{1}{8}e^{-x}\right).$$

The steady-state buffer empty probability for the SFM of N' is

$$\ell' = F'(0) = \left(0, \frac{1}{8}, \frac{1}{8}\right).$$

In Figure 11, the plots of the elements F_1, F_2, F'_2 of the steady-state fluid PDFs $F = (F_1, F_2)$ and $F' = (F'_1, F'_2, F'_3)$ for the SFMs of N and N' as functions of x are depicted. It is sufficient to consider the functions $F_1(x) = \frac{1}{2} - \frac{1}{2}e^{-x}$, $F_2(x) = \frac{1}{2} - \frac{1}{4}e^{-x}$, $F'_2(x) = \frac{1}{4} - \frac{1}{8}e^{-x}$ only, since $F_1 = F'_1$ and $F'_2 = F'_3$.

We have $(DRS(N) \cup DRS(N'))/\mathcal{R}_{ft}(N, N') = \{\mathcal{H}_1, \mathcal{H}_2\}$, where $\mathcal{H}_1 = \{M_1, M'_1\}$ and $\mathcal{H}_2 = \{M_2, M'_2, M'_3\}$.

First, consider the equivalence class \mathcal{H}_1 .




FIG. 11. The elements of the steady-state fluid PDFs for the SFMs of N and N' as functions of x

- The aggregate steady-state probabilities for \mathcal{H}_1 coincide: $\varphi_{\mathcal{H}_1 \cap DRS(N)} = \sum_{\{i|M_i \in \mathcal{H}_1 \cap DRS(N)\}} \varphi_i = \varphi_1 = \frac{1}{2} = \varphi'_1 = \sum_{\{j|M'_j \in \mathcal{H}_1 \cap DRS(N')\}} \varphi'_j = \varphi'_{\mathcal{H}_1 \cap DRS(N')}$.
- The aggregate steady-state buffer empty probabilities for \mathcal{H}_1 coincide: $\ell_{\mathcal{H}_1 \cap DRS(N)} = \sum_{\{i|M_i \in \mathcal{H}_1 \cap DRS(N)\}} \ell_i = \ell_1 = 0 = \ell'_1 = \sum_{\{j|M'_j \in \mathcal{H}_1 \cap DRS(N')\}} \ell'_j = \ell'_{\mathcal{H}_1 \cap DRS(N')}$.
- The aggregate steady-state fluid PDFs for \mathcal{H}_1 coincide: $F_{\mathcal{H}_1 \cap DRS(N)}(x) = \sum_{\{i|M_i \in \mathcal{H}_1 \cap DRS(N)\}} F_i(x) = F_1(x) = \frac{1}{2} - \frac{1}{2}e^{-x} = F'_1(x) = \sum_{\{j|M'_j \in \mathcal{H}_1 \cap DRS(N')\}} F'_j(x) = F'_{\mathcal{H}_1 \cap DRS(N')}(x)$, where $x > 0$.
- The aggregate steady-state fluid probability density functions for \mathcal{H}_1 coincide: $f_{\mathcal{H}_1 \cap DRS(N)}(x) = \sum_{\{i|M_i \in \mathcal{H}_1 \cap DRS(N)\}} f_i(x) = f_1(x) = \frac{1}{2}e^{-x} = f'_1(x) = \sum_{\{j|M'_j \in \mathcal{H}_1 \cap DRS(N')\}} f'_j(x) = f'_{\mathcal{H}_1 \cap DRS(N')}(x)$, where $x > 0$.

Second, consider the equivalence class \mathcal{H}_2 .

- The aggregate steady-state probabilities for \mathcal{H}_2 coincide: $\varphi_{\mathcal{H}_2 \cap DRS(N)} = \sum_{\{i|M_i \in \mathcal{H}_2 \cap DRS(N)\}} \varphi_i = \varphi_2 = \frac{1}{2} = \frac{1}{4} + \frac{1}{4} = \varphi'_2 + \varphi'_3 = \sum_{\{j|M'_j \in \mathcal{H}_2 \cap DRS(N')\}} \varphi'_j = \varphi'_{\mathcal{H}_2 \cap DRS(N')}$.
- The aggregate steady-state buffer empty probabilities for \mathcal{H}_2 coincide: $\ell_{\mathcal{H}_2 \cap DRS(N)} = \sum_{\{i|M_i \in \mathcal{H}_2 \cap DRS(N)\}} \ell_i = \ell_2 = \frac{1}{4} = \frac{1}{8} + \frac{1}{8} = \ell'_2 + \ell'_3 = \sum_{\{j|M'_j \in \mathcal{H}_2 \cap DRS(N')\}} \ell'_j = \ell'_{\mathcal{H}_2 \cap DRS(N')}$.
- The aggregate steady-state fluid PDFs for \mathcal{H}_2 coincide: $F_{\mathcal{H}_2 \cap DRS(N)}(x) = \sum_{\{i|M_i \in \mathcal{H}_2 \cap DRS(N)\}} F_i(x) = F_2(x) = \frac{1}{2} - \frac{1}{4}e^{-x} = \frac{1}{4} - \frac{1}{8}e^{-x} + \frac{1}{4} - \frac{1}{8}e^{-x} = F'_2(x) + F'_3(x) = \sum_{\{j|M'_j \in \mathcal{H}_2 \cap DRS(N')\}} F'_j(x) = F'_{\mathcal{H}_2 \cap DRS(N')}(x)$, where $x > 0$.
- The aggregate steady-state fluid probability density functions for \mathcal{H}_2 coincide: $f_{\mathcal{H}_2 \cap DRS(N)}(x) = \sum_{\{i|M_i \in \mathcal{H}_2 \cap DRS(N)\}} f_i(x) = f_2(x) = \frac{1}{4}e^{-x} = \frac{1}{8}e^{-x} + \frac{1}{8}e^{-x} = f'_2(x) + f'_3(x) = \sum_{\{j|M'_j \in \mathcal{H}_2 \cap DRS(N')\}} f'_j(x) = f'_{\mathcal{H}_2 \cap DRS(N')}(x)$, where $x > 0$.

One can also see that $\varphi_{\leftrightarrow f_1} = \varphi'_{\leftrightarrow f_1} = \varphi$, $\ell_{\leftrightarrow f_1} = \ell'_{\leftrightarrow f_1} = \ell$, $F_{\leftrightarrow f_1}(x) = F'_{\leftrightarrow f_1}(x) = F(x)$, $x > 0$, and $f_{\leftrightarrow f_1}(x) = f'_{\leftrightarrow f_1}(x) = f(x)$, $x > 0$.




FIG. 12. The diagram of the document preparation system

9. DOCUMENT PREPARATION SYSTEM

Let us consider an application example describing three different models of a document preparation system. The system receives (in an arbitrary order or in parallel) the collections of the text and graphics files as its inputs and writes them into the operative memory of a computer. The system then reads the (mixed) data from there and produces properly formatted output documents consisting of text and images. In general, it is supposed that the text file collections are transferred into the operative memory slower, but for longer time than the graphics ones. In detail, the low resolution graphics is transferred into the operative memory with the same speed as the high resolution one, but it takes less time than for the latter. The data from the operative memory is consumed for processing quicker, but for shorter time than the input file collections of any type. The operative memory capacity is supposed to be unlimited (for example, there exist some special mechanisms to ensure that the memory upper boundary can always be increased, such as using the page file, stored on a hard drive of the computer). The lower boundary of the operative memory is zero. The diagram of the system is depicted in Figure 12.

The meaning of the actions that label the transitions of the LFSPNs which will specify the three models of the document preparation system is as follows. The action tx represents writing the text files into the operative memory. The action gr represents putting the graphics files into the operative memory. Particularly, the action gl corresponds to writing the low resolution graphics while gh specifies writing the high resolution graphics. The action dt represents reading the data (consisting of the portions of the input text and images) from the operative memory. In each LFSPN, a single continuous place containing fluid will represent the operative memory with a data volume stored.

In Figure 13, the LFSPNs N and N' specifying the *standard* document preparation system, as well as the LFSPN N'' representing the *enhanced* one that differentiates between the low and high resolution graphics, are presented. The rate of all transitions labeled with the action tx is 1, the rate of those labeled with gr is 2 and the rate of those labeled with dt is 3. Further, the rate of the transition with the label gl is $\frac{3}{2}$ and the rate of that with the label gh is $\frac{1}{2}$. The rate of the fluid flow




FIG. 13. The LFSPNs of the standard and enhanced document preparation systems

along the continuous arcs from the transitions labeled with the action tx is 1 while that from the transitions labeled with gr is 2. Next, the fluid flow rate from the transitions with the label gl or gh is the same and equals 1. The rate of the fluid flow along the continuous arcs to the transitions labeled with the action dt is 7.

We have $N \xleftrightarrow{fl} N'$. Since LFSPNs have an *interleaving* semantics due to the *continuous* time approach and the *race* condition applied to transition firings, the parallel execution of actions (here in N) is modeled by the sequential non-determinism (in N'). Fluid bisimulation equivalence is an interleaving relation constructed in conformance with the LFSPNs semantics. In our application example, one can see that the “sequential” LFSPN N' may be replaced with the fluid bisimulation equivalent and structurally simpler “concurrent” LFSPN N , the latter having less transitions and arcs. Thus, the mentioned equivalence can be used not just to reduce behaviour of LFSPNs, but also to simplify their structure.

We have $DRS(N) = \{M_1, M_2, M_3, M_4\}$, where $M_1 = (1, 1, 0, 0)$, $M_2 = (0, 1, 1, 0)$, $M_3 = (1, 0, 0, 1)$, $M_4 = (0, 0, 1, 1)$; $DRS(N') = \{M'_1, M'_2, M'_3, M'_4\}$, where $M'_1 = (1, 0, 0, 0)$, $M'_2 = (0, 1, 0, 0)$, $M'_3 = (0, 0, 1, 0)$, $M'_4 = (0, 0, 0, 1)$; and $DRS(N'') = \{M''_1, M''_2, M''_3, M''_4, M''_5, M''_6\}$, where $M''_1 = (1, 1, 0, 0, 0)$, $M''_2 = (1, 0, 1, 0, 0)$, $M''_3 = (0, 1, 0, 0, 1)$, $M''_4 = (1, 0, 0, 1, 0)$, $M''_5 = (0, 0, 1, 0, 1)$, $M''_6 = (0, 0, 0, 1, 1)$.

In Figure 14, the discrete reachability graphs $DRG(N)$, $DRG(N')$, $DRG(N'')$ are depicted. The discrete parts of the LFSPNs N and N' have the same behaviour.

Let N''' is an abstraction of N'' by assuming that the actions gl and gh coincide with the action gr . Then it holds $N \xleftrightarrow{fl} N' \xleftrightarrow{fl} N'''$. In such a case, $DRS(N''') = \{M'''_1, M'''_2, M'''_3, M'''_4, M'''_5, M'''_6\}$ coincides with $DRS(N'')$ up to the trivial renaming bijection on the places. Further, $DRG(N''')$ coincides with $DRG(N'')$ up to the analogous renaming the transitions.

Let $\mathcal{K}_1 = \{M_1\}$, $\mathcal{K}_2 = \{M_2\}$, $\mathcal{K}_3 = \{M_3\}$, $\mathcal{K}_4 = \{M_4\}$ and $\mathcal{K}'_1 = \{M'_1\}$, $\mathcal{K}'_2 = \{M'_2\}$, $\mathcal{K}'_3 = \{M'_3\}$, $\mathcal{K}'_4 = \{M'_4\}$, as well as $\mathcal{K}'''_1 = \{M'''_1\}$, $\mathcal{K}'''_2 = \{M'''_2, M'''_4\}$, $\mathcal{K}'''_3 = \{M'''_3\}$, $\mathcal{K}'''_4 = \{M'''_5, M'''_6\}$. In Figure 15, the quotient (by \xleftrightarrow{fl}) discrete reachability




FIG. 14. The discrete reachability graphs of the LFSPNs of the standard and enhanced document preparation systems




FIG. 15. The quotient discrete reachability graphs of the LFSPNs of the standard document preparation system and that of the abstract LFSPN of the enhanced document preparation system

graphs $DRG_{\leftrightarrow_{fl}}(N)$, $DRG_{\leftrightarrow_{fl}}(N')$, $DRG_{\leftrightarrow_{fl}}(N''')$ are depicted. Obviously, $DRG_{\leftrightarrow_{fl}}(N) \simeq DRG_{\leftrightarrow_{fl}}(N') \simeq DRG_{\leftrightarrow_{fl}}(N''')$. Then it is clear that the discrete parts of the LFSPNs N , N' and N''' have the same *quotient* behaviour. Thus, quotienting by fluid bisimulation equivalence can be used to substantially reduce behaviour of LFSPNs. It is also clear that the discrete parts of the LFSPNs N and N' have the same *complete* and *quotient* behaviour.

The sojourn time average and variance vectors of N''' are

$$SJ''' = \left(\frac{1}{3}, 1, \frac{1}{2}, 1, \frac{1}{3}, \frac{1}{3} \right), \quad VAR''' = \left(\frac{1}{9}, 1, \frac{1}{4}, 1, \frac{1}{9}, \frac{1}{9} \right).$$

The complete and quotient sojourn time average and variance vectors of N and N' , as well as the quotient corresponding vectors of N''' , are

$$SJ = SJ_{\leftrightarrow_{fl}} = SJ' = SJ'_{\leftrightarrow_{fl}} = SJ'''_{\leftrightarrow_{fl}} = \left(\frac{1}{3}, \frac{1}{2}, 1, \frac{1}{3} \right),$$

$$VAR = VAR_{\leftrightarrow_{fl}} = VAR' = VAR'_{\leftrightarrow_{fl}} = VAR'''_{\leftrightarrow_{fl}} = \left(\frac{1}{9}, \frac{1}{4}, 1, \frac{1}{9} \right).$$

The TRM \mathbf{Q}''' for $CTMC(N''')$ and FRM \mathbf{R}''' for the SFM of N''' are

$$\mathbf{Q}''' = \begin{pmatrix} -3 & \frac{3}{2} & 1 & \frac{1}{2} & 0 & 0 \\ 0 & -1 & 0 & 0 & 1 & 0 \\ 0 & 0 & -2 & 0 & \frac{3}{2} & \frac{1}{2} \\ 0 & 0 & 0 & -1 & 0 & 1 \\ 3 & 0 & 0 & 0 & -3 & 0 \\ 3 & 0 & 0 & 0 & 0 & -3 \end{pmatrix}, \quad \mathbf{R}''' = \begin{pmatrix} 3 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -7 & 0 \\ 0 & 0 & 0 & 0 & 0 & -7 \end{pmatrix}.$$

The TRMs \mathbf{Q} , $\mathbf{Q}_{\leftrightarrow fl}$, \mathbf{Q}' , $\mathbf{Q}'_{\leftrightarrow fl}$, $\mathbf{Q}'''_{\leftrightarrow fl}$ for $CTMC(N)$, $CTMC_{\leftrightarrow fl}(N)$, $CTMC(N')$, $CTMC_{\leftrightarrow fl}(N')$, $CTMC_{\leftrightarrow fl}(N''')$, and FRMs \mathbf{R} , $\mathbf{R}_{\leftrightarrow fl}$, \mathbf{R}' , $\mathbf{R}'_{\leftrightarrow fl}$, $\mathbf{R}'''_{\leftrightarrow fl}$ for the complete and quotient SFMs of N , N' and quotient SFM of N''' are

$$\mathbf{Q} = \mathbf{Q}_{\leftrightarrow fl} = \mathbf{Q}' = \mathbf{Q}'_{\leftrightarrow fl} = \mathbf{Q}'''_{\leftrightarrow fl} = \begin{pmatrix} -3 & 1 & 2 & 0 \\ 0 & -2 & 0 & 2 \\ 0 & 0 & -1 & 1 \\ 3 & 0 & 0 & -3 \end{pmatrix},$$

$$\mathbf{R} = \mathbf{R}_{\leftrightarrow fl} = \mathbf{R}' = \mathbf{R}'_{\leftrightarrow fl} = \mathbf{R}'''_{\leftrightarrow fl} = \begin{pmatrix} 3 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -7 \end{pmatrix}.$$

Thus, the respective discrete and continuous parts of the LFSPNs N and N' have the same *complete* and *quotient* behaviour while N''' has the same *quotient* one. Therefore, it is enough to consider only LFSPN N from now on.

The discrete markings of LFSPN N are interpreted as follows: M_1 : both the text and graphics file collections are written to the memory, M_2 : the text file collection is resided in the memory and the graphics one is written to the memory, M_3 : the graphics file collection is resided in the memory and the text one is written to the memory, M_4 : the text and graphics file collections are resided in the memory and the data is read from there (if it is not empty).

We have $DRS^-(N) = \{M_4\}$, $DRS^0(N) = \emptyset$ and $DRS^+(N) = \{M_1, M_2, M_3\}$.

The steady-state PMF for $CTMC(N)$ is

$$\varphi = \left(\frac{2}{9}, \frac{1}{9}, \frac{4}{9}, \frac{2}{9} \right).$$

Then the *stability condition* for the SFM of N is fulfilled: $FluidFlow(q) = \sum_{i=1}^4 \varphi_i RP(M_i) = \frac{2}{9} \cdot 3 + \frac{1}{9} \cdot 2 + \frac{4}{9} \cdot 1 + \frac{2}{9}(-7) = -\frac{2}{9} < 0$.

For each eigenvalue γ we must have $|\gamma \mathbf{R} - \mathbf{Q}| =$

$$\begin{vmatrix} 3(\gamma+1) & -1 & -2 & 0 \\ 0 & 2(\gamma+1) & 0 & -2 \\ 0 & 0 & \gamma+1 & -1 \\ -3 & 0 & 0 & -7\gamma+3 \end{vmatrix} = -42\gamma^4 - 108\gamma^3 - 72\gamma^2 - 6\gamma = 0; \text{ hence,}$$

$$\gamma_1 = 0, \gamma_2 = -1, \gamma_3 = -\frac{1}{14}(11 + \sqrt{93}), \gamma_4 = -\frac{1}{14}(11 - \sqrt{93}).$$

The corresponding eigenvectors are the solutions of

$$\begin{aligned}
v_1 \begin{pmatrix} 3 & -1 & -2 & 0 \\ 0 & 2 & 0 & -2 \\ 0 & 0 & 1 & -1 \\ -3 & 0 & 0 & 3 \end{pmatrix} = 0, \quad v_2 \begin{pmatrix} 0 & -1 & -2 & 0 \\ 0 & 0 & 0 & -2 \\ 0 & 0 & 0 & -1 \\ -3 & 0 & 0 & 10 \end{pmatrix} = 0, \\
v_3 \begin{pmatrix} \frac{3}{14}(3-\sqrt{93}) & -1 & -2 & 0 \\ 0 & \frac{1}{7}(3-\sqrt{93}) & 0 & -2 \\ 0 & 0 & \frac{1}{14}(3-\sqrt{93}) & -1 \\ -3 & 0 & 0 & \frac{1}{2}(17+\sqrt{93}) \end{pmatrix} = 0, \\
v_4 \begin{pmatrix} \frac{3}{14}(3+\sqrt{93}) & -1 & -2 & 0 \\ 0 & \frac{1}{7}(3+\sqrt{93}) & 0 & -2 \\ 0 & 0 & \frac{1}{14}(3+\sqrt{93}) & -1 \\ -3 & 0 & 0 & \frac{1}{2}(17-\sqrt{93}) \end{pmatrix} = 0.
\end{aligned}$$

Then the eigenvectors are $v_1 = (\frac{2}{9}, \frac{1}{9}, \frac{4}{9}, \frac{2}{9})$, $v_2 = (0, -1, 2, 0)$,
 $v_3 = (\frac{14}{3-\sqrt{93}}, \frac{98}{(3-\sqrt{93})^2}, \frac{392}{(3-\sqrt{93})^2}, 1)$, $v_4 = (\frac{14}{3+\sqrt{93}}, \frac{98}{(3+\sqrt{93})^2}, \frac{392}{(3+\sqrt{93})^2}, 1)$.
Since $\varphi = F(\infty) = a_1 v_1$, we have $F(x) = \varphi + a_2 e^{72x} v_2 + a_3 e^{73x} v_3 + a_4 e^{74x} v_4$
and $a_1 = 1$. Since $\forall M_l \in DRS^+(N)$ $F_l(0) = \varphi_l + a_2 v_{2l} + a_3 v_{3l} + a_4 v_{4l} = 0$
and $DRS^+(N) = \{M_1, M_2, M_3\}$, we have the following linear equation system:

$$\begin{cases} \varphi_1 + a_2 v_{21} + a_3 v_{31} + a_4 v_{41} = \frac{2}{9} + \frac{14}{3-\sqrt{93}} a_3 + \frac{14}{3+\sqrt{93}} a_4 = 0 \\ \varphi_2 + a_2 v_{22} + a_3 v_{32} + a_4 v_{42} = \frac{1}{9} - a_2 + \frac{98}{(3-\sqrt{93})^2} a_3 + \frac{98}{(3+\sqrt{93})^2} a_4 = 0 \\ \varphi_3 + a_2 v_{23} + a_3 v_{33} + a_4 v_{43} = \frac{4}{9} + 2a_2 + \frac{392}{(3-\sqrt{93})^2} a_3 + \frac{392}{(3+\sqrt{93})^2} a_4 = 0 \end{cases}.$$

By solving the system, we get $a_2 = 0$, $a_3 = \frac{2(31-3\sqrt{93})}{93(3+\sqrt{93})}$, $a_4 = -\frac{2(10+\sqrt{93})}{21\sqrt{93}}$. Thus,
 $F(x) = (\frac{2}{9}, \frac{1}{9}, \frac{4}{9}, \frac{2}{9}) + \frac{2(31-3\sqrt{93})}{93(3+\sqrt{93})} e^{-\frac{1}{14}(11+\sqrt{93})x} (\frac{14}{3-\sqrt{93}}, \frac{98}{(3-\sqrt{93})^2}, \frac{392}{(3-\sqrt{93})^2}, 1) -$
 $\frac{2(10+\sqrt{93})}{21\sqrt{93}} e^{-\frac{1}{14}(11-\sqrt{93})x} (\frac{14}{3+\sqrt{93}}, \frac{98}{(3+\sqrt{93})^2}, \frac{392}{(3+\sqrt{93})^2}, 1)$.

Then the steady-state fluid PDF for the SFM of N is

$$\begin{aligned}
F(x) = & \left(\frac{2}{9} - \frac{(31-3\sqrt{93})}{279} e^{-\frac{1}{14}(11+\sqrt{93})x} - \frac{4(10+\sqrt{93})}{3\sqrt{93}(3+\sqrt{93})} e^{-\frac{1}{14}(11-\sqrt{93})x}, \right. \\
& \frac{1}{9} - \frac{7(31-3\sqrt{93})}{279(3-\sqrt{93})} e^{-\frac{1}{14}(11+\sqrt{93})x} - \frac{28(10+\sqrt{93})}{3\sqrt{93}(3+\sqrt{93})^2} e^{-\frac{1}{14}(11-\sqrt{93})x}, \\
& \frac{4}{9} - \frac{28(31-3\sqrt{93})}{279(3-\sqrt{93})} e^{-\frac{1}{14}(11+\sqrt{93})x} - \frac{112(10+\sqrt{93})}{3\sqrt{93}(3+\sqrt{93})^2} e^{-\frac{1}{14}(11-\sqrt{93})x}, \\
& \left. \frac{2}{9} + \frac{2(31-3\sqrt{93})}{93(3+\sqrt{93})} e^{-\frac{1}{14}(11+\sqrt{93})x} - \frac{2(10+\sqrt{93})}{21\sqrt{93}} e^{-\frac{1}{14}(11-\sqrt{93})x} \right).
\end{aligned}$$

The steady-state fluid probability density function for the SFM of N is

$$\begin{aligned}
f(x) = \frac{dF(x)}{dx} = & \left(\frac{e^{-\frac{1}{14}(11+\sqrt{93})x} (31-\sqrt{93}+(31+\sqrt{93})e^{\frac{\sqrt{93}x}{7}})}{1953}, \frac{e^{-\frac{1}{14}(11+\sqrt{93})x} (-1+e^{\frac{\sqrt{93}x}{7}})}{9\sqrt{93}}, \right. \\
& \left. \frac{4e^{-\frac{1}{14}(11+\sqrt{93})x} (-1+e^{\frac{\sqrt{93}x}{7}})}{9\sqrt{93}}, \frac{e^{-\frac{1}{14}(11+\sqrt{93})x} (14(-31+\sqrt{93})+(620+48\sqrt{93})e^{\frac{\sqrt{93}x}{7}})}{4557(3+\sqrt{93})} \right).
\end{aligned}$$

The steady-state buffer empty probability for the SFM of N is

$$\ell = F(0) = \left(0, 0, 0, \frac{2}{63} \right).$$

In Figure 16, the plots of the elements F_1, F_2, F_3, F_4 of the steady-state fluid PDF $F = (F_1, F_2, F_3, F_4)$ for the SFM of N , as functions of x , are depicted.




FIG. 16. The elements of the steady-state fluid PDF for the SFM of the concurrent LFSPN of the document preparation system

We can now calculate for the document preparation system some steady-state performance indices, based on the discrete and continuous measures from [53].

- The *fraction of time when both the text and graphics file collections are written to the memory*, calculated as the *fraction (proportion) of time spent in the set of discrete markings* $\{M_1\} \subseteq DRS(N)$, is

$$TimeFract(\{M_1\}) = \sum_{\{i|M_i \in \{M_1\}\}} \varphi_i = \varphi_1 = \frac{2}{9}.$$

- The *average number of the text file collections received per unit of time*, calculated as the *firing frequency (throughput) of the transition* $t_1 \in T_N$ (average number of firings per unit of time), is

$$FiringFreq(t_1) = \sum_{\{i|t_1 \in Ena(M_i), M_i \in DRS(N)\}} \varphi_i \Omega_N(t_1, M_i) = \varphi_1 \Omega_N(t_1, M_1) = \frac{2}{9} \cdot 1 = \frac{2}{9}.$$

- The *throughput of the system*, calculated as the *firing frequency (throughput) of the transition* $t_3 \in T_N$ (average number of firings per unit of time), is

$$FiringFreq(t_3) = \sum_{\{i|t_3 \in Ena(M_i), M_i \in DRS(N)\}} \varphi_i \Omega_N(t_3, M_i) = \varphi_4 \Omega_N(t_3, M_4) = \frac{2}{9} \cdot 3 = \frac{2}{3}.$$

- The *probability that the memory is not empty*, calculated as the *probability of a positive fluid level in a continuous place* $q \in Pc_N$, is

$$\begin{aligned}
FluidLevel(q) &= \sum_{\{i|M_i \in DRS(N)\}} \left(\ell_i \cdot 0 + \int_{0+}^{\infty} f_i(x) \cdot 1 dx \right) = \\
&= \sum_{\{i|M_i \in DRS(N)\}} \int_{0+}^{\infty} f_i(x) dx = \sum_{\{i|M_i \in DRS(N)\}} (\varphi_i - \ell_i) = \\
1 - \sum_{\{i|M_i \in DRS(N)\}} \ell_i &= 1 - (\ell_1 + \ell_2 + \ell_3 + \ell_4) = 1 - \frac{2}{63} = \frac{61}{63}.
\end{aligned}$$

- The probability that the operative memory contains at least 5 Mb data, calculated as the probability that the fluid level in a continuous place $q \in Pc_N$ does not lie below the value $5 \in \mathbb{R}_{>0}$, is

$$\begin{aligned}
FluidLevel(q, 5) &= \sum_{\{i|M_i \in DRS(N)\}} \left(\ell_i \cdot 0 + \int_{0+}^5 f_i(x) \cdot 0 dx + \int_5^{\infty} f_i(x) \cdot 1 dx \right) = \\
&= \sum_{\{i|M_i \in DRS(N)\}} \int_5^{\infty} f_i(x) dx = \sum_{\{i|M_i \in DRS(N)\}} (\varphi_i - F_i(5)) = \\
1 - \sum_{\{i|M_i \in DRS(N)\}} F_i(5) &= 1 - (F_1(5) + F_2(5) + F_3(5) + F_4(5)) = \\
&= \frac{e^{-\frac{5}{14}(11+\sqrt{93})}(5673-631\sqrt{93}) + e^{\frac{5\sqrt{93}}{7}}(5673+631\sqrt{93})}{11718} \approx 0.6181.
\end{aligned}$$

10. CONCLUSION

In this paper, we have defined two behavioural equivalences that preserve the qualitative and quantitative behavior of LFSPNs, related to both their discrete part (labeled CTSPNs and the underlying CTMCs) and continuous part (the associated SFMs). We have proposed on LFSPNs a linear-time relation of fluid trace equivalence and a branching-time relation of fluid bisimulation equivalence. Both equivalences respect *functional activity*, *stochastic timing* and *fluid flow*. We have demonstrated that fluid trace equivalence preserves average potential fluid change volume for the transition sequences of each given length. We have proven that fluid bisimulation equivalence strictly implies fluid trace one. We have explained how to reduce the discrete reachability graphs and underlying CTMCs of LFSPNs via quotienting the respective labeled transition systems by the largest fluid bisimulation. We have proven that fluid bisimulation equivalence preserves the qualitative and stationary quantitative behaviour, hence, the functionality and performance measures of the equivalent systems coincide. We have presented a case study of the three LFSPNs, all modeling the document preparation system, with a goal to show how fluid bisimulation equivalence can be used to simplify the LFSPNs structure and behaviour.

We plan to define a fluid place bisimulation relation that connects “similar” continuous places of LFSPNs, like place bisimulations [3, 2, 50, 51, 52] relate discrete places of (standard) Petri nets. The *lifting* of the relation to the discrete-continuous LFSPN markings (with discrete markings treated as the multisets of places) will respect both the fluid distribution among the related continuous places and the rates of fluid flow through them. For this purpose, we should introduce a novel notion of the multiset analogue with non-negative real-valued multiplicities of the elements. While multiset is a mapping from a countable set to all natural numbers, we need a more sophisticated mapping from the set of continuous places to all

non-negative real numbers, corresponding to the associated fluid levels. Such an extension of the multiset notion may use the results of [12, 49], concerning hybrid sets (the multiplicities of the elements are arbitrary integers) and fuzzy multisets (the multiplicities belong to the interval $[0;1]$). In this way, both the initial amount of fluid and its transit flow rate in each discrete marking may be distributed among several continuous places of an LFSPN, such that all of them are bisimilar to a particular continuous place of the equivalent LFSPN. The interesting point here is that fluid distributed among several bisimilar continuous places should be taken as the fluid contained in a single continuous place, resulting from aggregating those “constituent” continuous places with the use of fluid place bisimulation. Then the fluid level in the “aggregate” continuous place will be a sum of the fluid levels in the “constituent” continuous places. The probability density function for the sum of random variables representing the fluid levels in the “constituent” continuous places is defined via *convolution*. In this approach, we should avoid or treat correctly the situations when the fluid flow in the “aggregate” continuous place becomes suddenly non-continuous. This happens when some of the “constituent” continuous places are emptied while the others still contain a positive amount of fluid. Such a discontinuity is a result of applying the aggregation since it is not caused by either reaching the lower fluid boundary (zero fluid level) or change of the current discrete marking.

The summation of the fluid levels in the continuous places may be implemented with the constructions from [28] for *extended FSPNs* (EFSPNs). EFSPNs have special deterministic fluid jump arcs, used to transfer a deterministic amount of fluid between two continuous places via intermediate stochastic transitions connecting both places (deterministic fluid transfer). Analogously, random fluid jump arcs in EFSPNs are used to transfer a random amount of fluid from one continuous place to another (random fluid transfer). We can also use fluid transitions, mentioned in [28] as a direction for future development of FSPNs. Fluid transitions that transfer fluid from their input to their output continuous places are used to implement fluid volume conservation. If one of the input continuous places of a fluid transition becomes empty (i.e. the lower fluid boundary is reached) then the rate of the transition should change in a certain way. The continuous arcs between continuous places and fluid transitions may have multiplicities that change the fluid flow along the arcs according to a factor. Fluid transitions may be controlled by a discrete marking, using the guard functions associated with them or applying the inhibitor and test arcs, i.e. by the constructions that do not affect discrete markings.

Further, we intend to apply to LFSPNs an analogue of the effective reduction technique based on the place bisimulations of Petri nets [3, 2]. In this way, we shall merge several equivalent continuous places and, in some cases, the transitions between them. This should result in the significant reductions of LFSPNs. The number of continuous places in an LFSPN impacts drastically the complexity of its solution. The analytical solution is normally possible for just a few continuous places (or even only for one). In all other cases, when modeling realistic large and complex systems, we have to apply numerical techniques to solve systems of partial differential equations, or the method of simulation. Hence, the reduction of the number of continuous places accomplished with the place bisimulation merging appears to be even more important for LFSPNs than for Petri nets.

Moreover, we intend to provide fluid bisimulation equivalence with logical characterization via a fluid extension of Probabilistic Modal Logic (PML) [43] for probabilistic transitions systems. For this, we can apply the characterization of strong equivalence by the logic PML_μ [24, 25], a stochastic extension of PML to the stochastic process algebra PEPA [38]. In addition, the logical characterizations of fluid trace and bisimulation equivalences can be constructed using the modal logics HML_{MT_r} and HML_{MB} [8, 6], based on the well-known logic HML [37]. The pointed modal logics characterize, respectively, Markovian trace and bisimulation equivalences on sequential and concurrent Markovian process calculi SMPC and CMPC. We can apply the results of [9], where on (sequential) Markovian process calculus MPC, the logical characterizations of Markovian trace and bisimulation equivalences are given with the HML-based modal logics HML_{NPMT_r} and HML_{MB} , respectively.

REFERENCES

- [1] ANGIUS A., HORVÁTH A., HALAWANI S.M., BARUKAB O., AHMAD A.R., BALBO G. *Use of flow equivalent servers in the transient analysis of product form queueing networks*. *Lecture Notes in Computer Science* **9081**, p. 15–29, Springer, 2015.
- [2] AUTANT C., PFISTER W., SCHNOEBELEN PH. *Place bisimulations for the reduction of labeled Petri nets with silent moves*. *Proceedings of 6th International Conference on Computing and Information - 94 (ICCI'94)*, p. 230–246, Trent University, Canada, 1994, <http://www.lsv.ens-cachan.fr/Publis/PAPERS/PS/APS-icci94.ps>.
- [3] AUTANT C., SCHNOEBELEN PH. *Place bisimulations in Petri nets*. *Lecture Notes in Computer Science* **616**, p. 45–61, Springer, 1992.
- [4] BALBO G. *Introduction to stochastic Petri nets*. *Lecture Notes in Computer Science* **2090**, p. 84–155, Springer, 2001.
- [5] BALBO G. *Introduction to generalized stochastic Petri nets*. *Lecture Notes in Computer Science* **4486**, p. 83–131, Springer, 2007.
- [6] BERNARDO M. *A survey of Markovian behavioral equivalences*. *Lecture Notes in Computer Science* **4486**, p. 180–219, Springer, 2007, <http://www.sti.uniurb.it/bernardo/documents/sfm07pe.tuto.pdf>.
- [7] BERNARDO M. *Non-bisimulation-based Markovian behavioral equivalences*. *Journal of Logic and Algebraic Programming* **72**, p. 3–49, Elsevier, 2007, <http://www.sti.uniurb.it/bernardo/documents/jlap72.pdf>.
- [8] BERNARDO M., BOTTA S. *Modal logic characterization of Markovian testing and trace equivalences*. *Proceedings of 1st International Workshop on Logic, Models and Computer Science - 06 (LMCS'06)* (F. Corradini, C. Toffalori, eds.), Camerino, Italy, April 2006, *Electronic Notes in Theoretical Computer Science* **169**, p. 7–18, Elsevier, 2006, <http://www.sti.uniurb.it/bernardo/documents/lmcs2006.pdf>.
- [9] BERNARDO M., BOTTA S. *A survey of modal logics characterizing behavioural equivalences for non-deterministic and stochastic systems*. *Mathematical Structures in Computer Science* **18**, p. 29–55, Cambridge University Press, 2008, <http://www.sti.uniurb.it/bernardo/documents/mcscs18.pdf>.
- [10] BERNARDO M., DE NICOLA R., LORETI M. *A uniform framework for modeling non-deterministic, probabilistic, stochastic, or mixed processes and their behavioral equivalences*. *Information and Computation* **225**, p. 29–82, Elsevier, 2013, <http://www.sti.uniurb.it/bernardo/documents/ic225.pdf>.
- [11] BERNARDO M., TESEI L. *Encoding timed models as uniform labeled transition systems*. *Lecture Notes in Computer Science* **8168**, p. 104–118, Springer, 2013.
- [12] BLIZARD W.D. *The development of multiset theory*. *The Review of Modern Logic* **1(4)**, p. 319–352, 1991, http://projecteuclid.org/download/pdf_1/euclid.rml/1204834739.
- [13] BOBBIO A., GARG S., GRIBAUDO M., HORVÁTH A., SERENO M., TELEK M. *Modeling software systems with rejuvenation, restoration and checkpointing through fluid stochastic Petri nets*. *Proceedings of 8th International Workshop on Petri Nets and Performance Models - 99 (PNPM'99)*, Zaragoza, Spain, p. 82–91, IEEE Computer Society Press, September 1999,

- <http://www.di.unito.it/~marcog/Downloads/PNPM99-BGGHST.pdf>,
<http://webspn.hit.bmc.hu/~telek/cikkekbobb99a.ps.gz>.
- [14] BOBBIO A., PULIAFITO A., TELEK M., TRIVEDI K.S. *Recent developments in non-Markovian stochastic Petri nets*. *Journal of Circuits, Systems and Computers* **8(1)**, p. 119–158, World Scientific, 1998, <http://webspn.hit.bmc.hu/~telek/cikkekbobb98c.ps.gz>,
http://people.ee.duke.edu/~kst/spn_papers/bobb98c.ps.
- [15] BUCHHOLZ P. *A notion of equivalence for stochastic Petri nets*. *Lecture Notes in Computer Science* **935**, p. 161–180, Springer, 1995.
- [16] BUCHHOLZ P. *Iterative decomposition and aggregation of labeled GSPNs*. *Lecture Notes in Computer Science* **1420**, p. 226–245, Springer, 1998.
- [17] CARDELLI L., TRIBASTONE M., TSCHAIKOWSKI M., VANDIN A. *Forward and backward bisimulations for chemical reaction networks*. *Proceedings of 26th International Conference on Concurrency Theory - 15 (CONCUR'15)*, Madrid, Spain, September 2015, *Leibniz International Proceedings in Informatics (LIPIcs)* **42**, p. 226–239, Dagstuhl Publishing, Leibniz-Zentrum für Informatik, Schloss Dagstuhl, Germany, 2015, <http://drops.dagstuhl.de/opus/volltexte/2015/5368/pdf/8.pdf>, <https://dl.dropboxusercontent.com/u/13100903/papers/concur2015.pdf>.
- [18] CARDELLI L., TRIBASTONE M., TSCHAIKOWSKI M., VANDIN A. *Comparing chemical reaction networks: a categorical and algorithmic perspective*. *Proceedings of 31st Annual ACM/IEEE Symposium on Logic in Computer Science - 16 (LICS'16)*, New York, USA, July 2016, 13 p., ACM Press, 2016, <https://dl.dropboxusercontent.com/u/13100903/papers/lics2016.pdf>.
- [19] CARDELLI L., TRIBASTONE M., TSCHAIKOWSKI M., VANDIN A. *Symbolic computation of differential equivalences*. *Proceedings of 43rd Annual ACM SIGPLAN-SIGACT Symposium on Principles of Programming Languages - 16 (POPL'16)*, St. Petersburg, Florida, USA, p. 137–150, ACM Press, January 2016, <http://sysma.imtlucca.it/wp-content/uploads/2015/11/z3-popl16.pdf>.
- [20] CARDELLI L., TRIBASTONE M., TSCHAIKOWSKI M., VANDIN A. *Efficient syntax-driven lumping of differential equations*. *Lecture Notes in Computer Science* **9636**, p. 93–111, Springer, 2016, <http://dl.dropboxusercontent.com/u/13100903/papers/tacas2016.pdf>.
- [21] CIARDO G., MUPPALA J.K., TRIVEDI K.S. *On the solution of GSPN reward models*. *Performance Evaluation* **12(4)**, p. 237–253, Elsevier, 1991, http://people.ee.duke.edu/~kst/spn_papers/gspnrew.ps.
- [22] CIARDO G., NICOL D., TRIVEDI K.S. *Discrete-event simulation of fluid stochastic Petri nets*. *Report 97-24*, 15 p., Institute for Computer Applications in Science and Engineering (ICASE), NASA Langley Research Center, Hampton, Virginia, USA, May 1997, <http://historical.ncstrl.org/tr/pdf/icase/TR-97-24.pdf>.
- [23] CIARDO G., NICOL D., TRIVEDI K.S. *Discrete-event simulation of fluid stochastic Petri nets*. *IEEE Transactions on Software Engineering* **25(2)**, p. 207–217, IEEE Computer Society Press, 1999, http://people.ee.duke.edu/~kst/spn_papers/Discrete.pdf.
- [24] CLARK G., GIMORE S., HILLSTON J. *Specifying performance measures for PEPA*. *Lecture Notes in Computer Science* **1601**, p. 211–227, Springer, 1999, <http://homepages.inf.ed.ac.uk/stg/publications/arts1999.pdf>.
- [25] CLARK G., GILMORE S., HILLSTON J., RIBAUDO M. *Exploiting modal logic to express performance measures*. *Lecture Notes in Computer Science* **1786**, p. 247–261, Springer, 2000, <http://homepages.inf.ed.ac.uk/stg/publications/tools2000.pdf>.
- [26] DERISAVI S., HERMANN S., SANDERS W.H. *Optimal state-space lumping of Markov chains*. *Information Processing Letters* **87(6)**, p. 309–315, Elsevier, 2003, http://www.perform.illinois.edu/Papers/USAN_papers/02DER01.pdf.
- [27] ELWALID A.I., MITRA D. *Statistical multiplexing with loss priorities in rate-based congestion control of high-speed networks*. *IEEE Transactions on Communications* **42(11)**, p. 2989–3002, IEEE Computer Society Press, 1994.
- [28] GRIBAUDO M. *Hybrid formalism for performance evaluation: theory and applications*. *Ph.D. thesis*, 198 p., Department of Computer Science, University of Turin, Turin, Italy, 2002, <http://www.di.unito.it/~marcog/Downloads/PhDThesis.pdf>.

- [29] GRIBAUDO M., HORVÁTH A. *Fluid stochastic Petri nets augmented with flush-out arcs: a transient analysis technique*. *IEEE Transactions on Software Engineering* **28**(10), p. 944–955, IEEE Computer Society Press, 2002, <http://www.di.unito.it/~marcog/Downloads/GrHo02.pdf>.
- [30] GRIBAUDO M., MANINI D., SERICOLA B., TELEK M. *Second order fluid models with general boundary behaviour*. *Annals of Operations Research* **160**, p. 69–82, Springer, 2008.
- [31] GRIBAUDO M., SERENO M. *Simulation of fluid stochastic Petri nets*. *Proceedings of 8th International Symposium on Modeling, Analysis and Simulation of Computer and Telecommunication Systems - 00 (MASCOTS'00)*, p. 231–239, San Francisco, USA, August 2000, <http://www.di.unito.it/~marcog/Downloads/MASCOTS00.pdf>.
- [32] GRIBAUDO M., SERENO M., HORVÁTH A., BOBBIO A. *Fluid stochastic Petri nets augmented with flush-out arcs: modelling and analysis*. *Discrete Event Dynamic Systems: Theory and Applications* **11**(1–2), p. 97–117, Springer, 2001, <http://www.di.unito.it/~marcog/Downloads/DEDS2001.pdf>, <http://www.mfn.unipmn.it/~bobbio/BIBLIO/PAPERS/ANN001/deds2001.pdf>.
- [33] GRIBAUDO M., TELEK M. *Fluid models in performance analysis*. *Lecture Notes in Computer Science* **4486**, p. 271–317, Springer, 2007, <http://webspn.hit.bme.hu/~telek/cikkek/grib07b.pdf>.
- [34] GRIBAUDO M., TELEK M. *Stationary analysis of fluid level dependent bounded fluid models*. *Performance Evaluation* **65**(3–4), p. 241–261, Elsevier, 2008, <http://webspn.hit.bme.hu/~telek/cikkek/grib07a.pdf>.
- [35] HAVERKORT B.R. *Markovian models for performance and dependability evaluation*. *Lecture Notes in Computer Science* **2090**, p. 38–83, Springer, 2001, <http://www-i2.informatik.rwth-aachen.de/Teaching/Seminar/VOSS2005/have01.pdf>.
- [36] HAYDEN R.A., BRADLEY J.T. *A fluid analysis framework for a Markovian process algebra*. *Theoretical Computer Science* **411**, p. 2260–2297, Elsevier, 2010.
- [37] HENNESSY M.C.B., MILNER R.A.J. *Algebraic laws for non-determinism and concurrency*. *Journal of the ACM* **32**(1), p. 137–161, ACM Press, 1985, <http://www.scss.tcd.ie/Matthew.Hennessy/pubs/old/HMjacm85.pdf>.
- [38] HILLSTON J. *A compositional approach to performance modelling*. 158 p., Cambridge University Press, UK, 1996, <http://www.dcs.ed.ac.uk/pepa/book.pdf>.
- [39] HORTON G., KULKARNI V.G., NICOL D.M., TRIVEDI K.S. *Fluid stochastic Petri nets: theory, applications, and solution techniques*. *European Journal of Operations Research* **105**(1), p. 184–201, Elsevier, 1998, http://people.ee.duke.edu/~kst/spn_papers/Fluid.pdf.
- [40] HORVÁTH A., GRIBAUDO M. *Matrix geometric solution of fluid stochastic Petri nets*. *Proceedings of 4th International Conference on Matrix-Analytic Methods in Stochastic Models - 02*, p. 163–182, Adelaide, Australia, July 2002, <http://www.di.unito.it/~horvath/publications/papers/HoGr02.ps>, <http://www.di.unito.it/~marcog/Downloads/HoGr02.pdf>.
- [41] IACOBELLI G., TRIBASTONE M., VANDIN A. *Differential bisimulation for a Markovian process algebra*. *Lecture Notes in Computer Science* **9234**, p. 293–306, Springer, 2015.
- [42] KULKARNI V.G. *Modeling and analysis of stochastic systems. Second edition*. *Texts in Statistical Science* **84**, 563 p., Chapman and Hall / CRC Press, 2009.
- [43] LARSEN K.G., SKOU A. *Bisimulation through probabilistic testing*. *Information and Computation* **94**(1), p. 1–28, Elsevier, 1991.
- [44] MARSAN M.A. *Stochastic Petri nets: an elementary introduction*. *Lecture Notes in Computer Science* **424**, p. 1–29, Springer, 1990.
- [45] MARSAN M.A., BALBO G., CONTE G., DONATELLI S., FRANCESCHINIS G. *Modelling with generalised stochastic Petri nets*. *Wiley Series in Parallel Computing*, John Wiley and Sons, 316 p., 1995, <http://www.di.unito.it/~greatspn/GSPN-Wiley/>.
- [46] MOLLOY M.K. *Performance analysis using stochastic Petri nets*. *IEEE Transactions on Computing* **31**(9), p. 913–917, IEEE Computer Society Press, 1982.
- [47] MOLLOY M.K. *Discrete time stochastic Petri nets*. *IEEE Transactions on Software Engineering* **11**(4), p. 417–423, IEEE Computer Society Press, 1985.
- [48] NATKIN S.O. *Les reseaux de Petri stochastiques et leur application a l'evaluation des systemes informatiques*. *Ph. D. thesis*, Conseratoire National des Arts et Metiers, France, 1980.
- [49] SYROPOULOS A. *Mathematics of multisetts*. *Lecture Notes in Computer Science* **2235**, p. 347–358, Springer, 2001.

- [50] TARASYUK I.V. *Place bisimulation equivalences for design of concurrent and sequential systems*. *Electronic Notes in Theoretical Computer Science* **18**, p. 191–206, Elsevier, 1998, <http://itar.iis.nsk.su/files/itar/pages/equpltcs.pdf>.
- [51] TARASYUK I.V. *τ -equivalences and refinement for Petri nets based design*. *Technische Berichte TUD-FI00-11*, 41 p., Fakultät Informatik, Technische Universität Dresden, Germany, November 2000, http://itar.iis.nsk.su/files/itar/pages/eqtbrtud_cov.pdf, http://www.qucosa.de/fileadmin/data/qucosa/documents/10037/tud_TB_2000-11.pdf.
- [52] TARASYUK I.V. *Equivalences for behavioural analysis of concurrent and distributed computing systems*. 321 p., Academic Publisher “Geo”, Novosibirsk, 2007 (ISBN 978-5-9747-0098-9, in Russian).
- [53] TARASYUK I.V., BUCHHOLZ P. *Bisimulation for fluid stochastic Petri nets*. *Bulletin of the Novosibirsk Computing Center, Series Computer Science, IIS Special Issue* **38**, p. 121–149, NCC Publisher, Novosibirsk, 2015 (ISSN 1680-6972), <http://itar.iis.nsk.su/files/itar/pages/fspneqnc.pdf>, <http://bulletin.iis.nsk.su/files/article/fspneqnc.pdf>.
- [54] TÓTH J., LI G., RABITZ H., TOMLIN A.S. *The effect of lumping and expanding on kinetic differential equations*. *SIAM Journal of Applied Mathematics* **57(6)**, p. 1531–1556, Society for Industrial and Applied Mathematics, 1997.
- [55] TRIVEDI K.S., KULKARNI V.G. *FSPNs: fluid stochastic Petri nets*. *Lecture Notes in Computer Science* **691**, p. 24–31, Springer, 1993, http://www.ee.duke.edu/~kst/spn_papers/pn93.ps.
- [56] TSCHAIKOWSKI M., TRIBASTONE M. *Exact fluid lumpability for Markovian process algebra*. *Lecture Notes in Computer Science* **7454**, p. 380–394, Springer, 2012.
- [57] TSCHAIKOWSKI M., TRIBASTONE M. *Tackling continuous state-space explosion in a Markovian process algebra*. *Theoretical Computer Science* **517**, p. 1–33, Elsevier, 2014.
- [58] TSCHAIKOWSKI M., TRIBASTONE M. *Exact fluid lumpability in Markovian process algebra*. *Theoretical Computer Science* **538**, p. 140–166, Elsevier, 2014.
- [59] TSCHAIKOWSKI M., TRIBASTONE M. *Extended differential aggregations in process algebra for performance and biology*. *Proceedings of 12th International Workshop on Quantitative Aspects of Programming Languages and Systems - 14 (QAPL’14)*, Grenoble, France, April 2014, *Electronic Proceedings in Theoretical Computer Science* **154**, p. 34–47, Open Publishing Association, 2014, http://eprints.imtlucca.it/2591/1/EPTCS_Tribastone_Tschaikowski.pdf, <http://arxiv.org/pdf/1406.2067v1.pdf>.
- [60] TSCHAIKOWSKI M., TRIBASTONE M. *A unified framework for differential aggregations in Markovian process algebra*. *Journal of Logical and Algebraic Methods in Programming* **84**, p. 238–258, Elsevier, 2015.
- [61] TSCHAIKOWSKI M., TRIBASTONE M. *Approximate reduction of heterogenous nonlinear models with differential hulls*. *IEEE Transactions on Automatic Control* **61(4)**, p. 1099–1104, IEEE Computer Society Press, April 2016, http://eprints.imtlucca.it/2730/1/tac_differential_hull.pdf.
- [62] WOLF V., BAIER C., MAJSTER-CEDERBAUM M. *Trace machines for observing continuous-time Markov chains*. *Proceedings of 3rd International Workshop on Quantitative Aspects of Programming Languages - 05 (QAPL’05)*, Edinburgh, UK, 2005, *Electronic Notes in Theoretical Computer Science* **153(2)**, p. 259–277, Elsevier, 2006.
- [63] WOLTER K. *Second order fluid stochastic Petri nets: an extension of GSPNs for approximate and continuous modelling*. *Proceedings of Workshop on Analytical and Numerical Modelling Techniques, 1st World Congress on Systems Simulation - 97*, p. 328–332, Singapore, September 1997, <http://page.mi.fu-berlin.de/katinkaw/publications/anmt.ps>.

IGOR VALERIEVICH TARASYUK
 A.P. ERSHOV INSTITUTE OF INFORMATICS SYSTEMS,
 SIBERIAN BRANCH OF THE RUSSIAN ACADEMY OF SCIENCES,
 ACAD. LAVRENTIEV PR. 6,
 630090 NOVOSIBIRSK, RUSSIAN FEDERATION
 Email address: itar@iis.nsk.su

PETER BUCHHOLZ
FACULTY OF COMPUTER SCIENCE,
TECHNICAL UNIVERSITY OF DORTMUND,
OTTO-HAHN-STR. 16,
44227 DORTMUND, GERMANY
Email address: `peter.buchholz@cs.uni-dortmund.de`